

Available online at www.sciencerepository.org

Science Repository



Case Report

A Febrile Return from Mecca: is it the obvious?

Ramzi Kurd^{1,2}, Yael M. Szterenlicht², Amos M. Yinnon^{2*} and Reuven Friedmann¹

¹Departments of Geriatrics Shaare Zedek Medical Center, affiliated with the Hebrew-University-Hadassah Medical School, Jerusalem, Israel

²Internal Medicine, Shaare Zedek Medical Center, affiliated with the Hebrew-University-Hadassah Medical School, Jerusalem, Israel

ARTICLE INFO

Article history:

Received 20 December, 2018

Accepted 16 January, 2019

Published 30 January, 2019

Keywords:

Middle East respiratory Syndroom (MERS)

MERS-CoV (Coronavirus)

Adult Respiratory Syndrome (ARDS)

Isolation procedures

ABSTRACT

We describe a 53-year-old female patient with diabetes mellitus and mild hypertension who made the ritual Hajj pilgrimage to Mecca. On the return trip to Jerusalem, she developed a respiratory illness, high fever and chills. We describe the initial and subsequent management of this patient, epidemiological aspects and the surprising final diagnosis. The take-home messages are: first, emergency department staff should demonstrate a high index of suspicion and put patients in isolation who meet specific case definition criteria. Second, even if a certain disease, in this instance MERS-CoV is the obvious diagnosis, open-mindedness and a differential diagnosis are always necessary in order to avoid misdiagnosis.

© 2018 Amos M. Yinnon. Hosting by Science Repository.

Introduction

Patients with fever and acute respiratory symptoms with a recent travel history to Saudia Arabia should be considered to have Middle East Respiratory Syndrome (MERS) [1]. The causative pathogen MERS-CoV (Coronavirus) was first described in 2012. The disease is endemic in dromedary camels of the Middle East and these are evidently the only animals responsible for transmission to human contacts [2]. Stored dromedary camel sera suggest that the virus was present in these camels several decades ago [3]. At the end of December 2018, there have been a total of 2279 laboratory-confirmed cases of MERS-CoV (Coronavirus) infection, with 806 associated deaths, a case fatality rate of 35.3% [4]. The majority of these patients were reported from Saudi Arabia: 1901 cases, including 732 related deaths 38.5%. Because of the high rate of transmission in medical settings, most hospitals have adopted official case definitions of MERS-CoV and isolate travelers from endemic countries meeting these definitions until laboratory tests have ruled out this infection [5].

Case presentation

A 53-year-old female patient with diabetes mellitus and mild hypertension made the ritual Hajj pilgrimage to Mecca, carefully continuing the medications prescribed for these conditions. On the return trip to Jerusalem, she developed weakness, muscle aches, a runny nose, sore throat, and mild cough. After 2-3 mildly symptomatic days, she developed a 39°C fever and chills, general weakness, a harsh cough and shortness of breath. The next day, the 13th day after her pilgrimage, she presented at our hospital's emergency department. The physical examination revealed tachycardia and tachypnea, an O₂ saturation of 85% at room air, her blood pressure was 100/64, and there were right lower lobe crackles. The peripheral blood count was 19,000/mm³ with 85% neutrophils; blood chemistry included glucose 268 mg/dl, HgBA1c was not available, sodium 134 meq/L, potassium 4.0 meq/L, and creatinine was 0.55 mg/dl BUN 12 mg/dl. Markers of inflammation included CRP of 3 mg/dl (normal 0-0.5) and lactate was 5.4 mmol/L.

*Correspondence to: Amos M. Yinnon, MD, Professor of Medicine, Shaare Zedek Medical Center, P.O. Box 3235, Jerusalem 91031, Israel; Tel: +972-2-6555076; Fax: +972-2-6666840; E-mail: yinnon@szmc.org.il

(normal 0.7–2.1). The chest radiogram showed large bilateral opacities, suggestive of pulmonary edema. The electrocardiogram showed ST-T changes in the antero-lateral leads and serum troponin was 0.44 ng/L (normal 0.04–0.3); 3 hours later troponin increased to 0.99. The patient was admitted to the coronary care unit with a diagnosis of non-ST elevation myocardial infarction and secondary pulmonary edema. An infectious disease specialist was consulted, who immediately raised the possibility of MERS-CoV infection. Blood, sputum, and urine cultures were obtained, in addition to viral serology and nasal culture for viruses. The patient was started on ceftriaxone, azithromycin and oseltamivir, and she was transferred to a high-efficiency particulate air (HEPA) filtered isolation room in the medical department. Revision of the admission chest radiogram led to suspicion of adult respiratory distress syndrome (ARDS).

Within 45 minutes after admission to the medical department the patient collapsed. She was successfully resuscitated and intubated and transferred to the intensive care unit; follow-up blood tests showed severe leukocytosis of 34,500 with 85% neutrophils, acute kidney injury with creatinine 1.36 mg/dl, BUN 21 mg/dl, lactate 6.13 mmol/L and her liver enzymes showed signs of liver shock. The patient died 12 hours after her admission to the intensive care unit. Cultures revealed a surprising abundance of clues regarding this unfortunate patient's final illness. First, the nasal culture was positive for Respiratory Syncytial Virus (RSV). Second, the sputum culture was positive for *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Streptococcus pneumoniae*. Third, two blood cultures were positive for *Streptococcus agalactiae* (*Streptococcus* group B, GBS). Finally, serology and PCR were negative for influenza and MERS CoV.

The probable cause of death was post-RSV bacterial pneumonia and sepsis due to *Streptococcus agalactiae*, complicated by ARDS and septic shock.

Discussion

Our patient developed a febrile, respiratory tract illness on the return trip from the Hajj to Mecca. She met case definition criteria of MERS-CoV and should have immediately been placed in isolation; however, only after several hours in hospital was, she transferred to a HEPA-filtered isolation room [5, 6]. If she had been confirmed ill with MERS-CoV quite likely a secondary outbreak with disastrous consequences could have occurred as has been described extensively from Korea and elsewhere [5, 7, 8].

Hajj associated infections include mainly minor viral, respiratory illnesses, including influenza and RSV, but also vaccine-preventable meningitis and MERS-CoV.¹ Our patient most probably had a non-specific viral infection, quite likely RSV, with a secondary bacterial infection due to *Streptococcus agalactiae* (Group B *Streptococcus*, GBS). The latter organism - although often associated with peri-natal infection - due to prophylactic antibiotics given to parturient women previously shown to be GBS carriers, currently is more often encountered in medical departments [9, 10]. Post viral bacterial infection is most commonly due to *Streptococcus pneumoniae* and *Staphylococcus aureus*; a PubMed search did not detect any description of *Streptococcus agalactiae* pneumonia or sepsis in the aftermath of a

respiratory tract infection [11]. Diabetes mellitus, malignancy, HIV infection, and advanced hepatic and renal disease have been identified as risk factors for invasive GBS infection [12]. GBS associated pneumonia or sepsis in these patients carry a case fatality rate in excess of 40%.

The presumed initial diagnosis of this patient was non-ST elevation myocardial infarction and secondary pulmonary edema. Only subsequently did the combination of clinical and laboratory factors lead to the consideration of possible pneumonia and sepsis - with initiation of antibiotic treatment and vasopressor support. The international multicenter prevalence study on sepsis (the IMPReSS study) has shown that adherence with all components of the surviving sepsis bundle varies between hospitals and countries, with associated varying rates of survival [13, 14]. Accordingly, early diagnosis and institution of appropriate treatment is essential.

In conclusion, emergency department staff should demonstrate a high index of suspicion and put patients in isolation who meet certain case definition criteria. Even if a certain disease, in this instance MERS-CoV is the obvious diagnosis, open-mindedness and a differential diagnosis are always necessary in order to avoid misdiagnosis.

Funding

This manuscript was prepared without specific funding.

REFERENCES

- Gautret P, Benkouiten S, Al-Tawfiq JA, Memish ZA (2016) Hajj-associated respiratory infections. *Travel Med Infect Dis* 20: 1-18. [[Crossref](#)]
- Reusken CB, Raj VS, Koopmans MP, Haagmans BL (2016) Cross host transmission in the emergence of MERS coronavirus. *Curr Opin Virol* 16: 55-62. [[Crossref](#)]
- Mohd HA, Al-Tawfiq JA, Memish ZA (2016) Middle East Respiratory Syndrome Coronavirus (MERS-CoV) origin and animal reservoir. *Virol J* 13: 87. [[Crossref](#)]
- EMRO Situation Report Dec 2018.
- Kim JY, Song JY, Yoon YK, Choi SH, Song YG, et al. (2015) Middle East respiratory syndrome infection control and prevention guideline for health care facilities. *Infect Chemother* 47: 278-302. [[Crossref](#)]
- Al-Dorzi HM, Alsolamy S, Arabi YM (2016) Critically ill patients with Middle East respiratory syndrome coronavirus infection. *Crit Care* 20: 65. [[Crossref](#)]
- Kim KH, Tandil TE, Choi JW, Moon JM, Kim MS (2017) Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak in South Korea, 2015: epidemiology, characteristics and public health implications. *J Hosp Infect* 95: 207-213. [[Crossref](#)]
- Cho SY, Kang JM, Ha YE, Park GE, Lee JY, et al. (2016) MERS-CoV outbreak following a single patient exposure in an emergency room in South Korea: an epidemiological outbreak study. *Lancet*. 388: 994-1001. [[Crossref](#)]
- Ballard MS, Schønheyder HC, Knudsen JD, Lyytikäinen O, Dryden M, et al. (2016) for the International Bacteremia Surveillance Collaborative. The changing epidemiology of group B streptococcus bloodstream infection: a multi-national population-based assessment. *Infect Dis (Lond)* 48: 386-391. [[Crossref](#)]
- Lamagni TL, Keshishian C, Efstratiou A, Guy R, Henderson KL, et al. (2013) Emerging trends in the epidemiology of invasive group B

- streptococcal disease in England and Wales, 1991-2010. *Clin Infect Dis* 57: 682-688. [[Crossref](#)]
11. Florescu DF, Kalil AC (2014) The complex link between influenza and severe sepsis. *Virulence* 5: 137-142. [[Crossref](#)]
 12. Cebey-López M, Pardo-Seco J, Gómez-Carballa A, Martínón-Torres N, Martínón-Sánchez JM, et al. (2016) Bacteremia in Children Hospitalized with Respiratory Syncytial Virus Infection. *PLoS One*. [[Crossref](#)]
 13. Rhodes A, Phillips G, Beale R, Cecconi M, Chiche JD, et al. (2015) The Surviving Sepsis Campaign bundles and outcome: results from the International Multicentre Prevalence Study on Sepsis (the IMPReSS study). *Intensive Care Med* 41: 1620-1628. [[Crossref](#)]
 14. Levy MM, Evans LE, Rhodes A (2018) The Surviving Sepsis Campaign Bundle: 2018 update. *Intensive Care Med* 44: 925-928. [[Crossref](#)]