

Available online at [www.sciencerepository.org](http://www.sciencerepository.org)

Science Repository



## Research Article

# Image-Guided Percutaneous Biopsy for the Diagnosis of Musculoskeletal Tumors: An Analysis of 698 Cases and a Literature Review

Giulia Trovarelli, Adriana Ivette Castaneda-Martinez, Elisa Pala, Alessandro Cappellari, Andrea Angelini, Antonio Berizzi and Pietro Ruggieri\*

Department of Orthopaedics and Orthopaedic Oncology, University of Padova, Padova, Italy

## ARTICLE INFO

## Article history:

Received: 13 May, 2020

Accepted: 29 May, 2020

Published: 30 June, 2020

## Keywords:

Core-needle

tru-cut needle

bone tumor

soft tissue tumor

diagnosis

imaging-guided biopsy

## ABSTRACT

**Purpose:** Incisional biopsy is still used to obtain material for a straightforward histological diagnosis in musculoskeletal lesions. The aim of our study was to evaluate 1) the percentage of diagnostic procedures, 2) diagnostic accuracy, and 3) the incidence of complications after imaging-guided percutaneous biopsy (PCB).

**Case Series:** This is a retrospective analysis of imaging-guided PCB performed between January 2016 and September 2019 by fluoroscopy in bone lesions or under ultrasound-guidance in soft tissue lesions. Specimens were classified as diagnostic or non-diagnostic according to the pathologist's report; diagnostic accuracy was determined by comparing histopathological results from biopsy and tumor resection. PCB was diagnostic in 94% and 97% of cases, with a diagnostic accuracy for detecting tumor disease in soft tissue and bone lesions of 98% and 100%, respectively. No complications were observed.

**Discussion:** The PCB is safe, minimally invasive, and cost-effective; thus, it should be the gold standard for the diagnosis of musculoskeletal lesions.

© 2020 Pietro Ruggieri. Hosting by Science Repository.

## Introduction

Diagnosis for patients with musculoskeletal tumors is based on history, clinical, radiological, and histological examination. Most frequent mistakes in treatment are due to delay in diagnosis or misdiagnosis, affecting the patient's survival and the possibility of limb salvage [1, 2]. When clinical and radiologic features do not allow to distinguish with certainty about biological behaviour or in patients with a history of malignancy, histological diagnosis is fundamental [1-6]. It is even more critical in sites in which salvage surgery is demanding, such as pelvis or limb reconstructions, where it is essential to preserve as much tissue as possible to allow soft tissue coverage and function [7-10]. Consequently, it is mandatory to always perform a biopsy before treating musculoskeletal lesions, with very few exceptions.

The biopsy is the last step of staging, and it is a compromise between the need to have significant tissue and the need to avoid local or systemic contamination [3-6]. Different procedures such as incisional biopsy (IB), fine-needle aspiration (FNA), or percutaneous biopsy (PCB) were used with specific advantages and disadvantages [11-44]. IB has high diagnostic accuracy but also complications up to 17% [11-14, 21, 24]. FNA has limited application in musculoskeletal oncology because of the need for architectural evaluation: it should only be used for diagnosis of recurrence [12-18]. PCB has few complications (less than 7%), even if its diagnostic accuracy is lower than IB (range, 76-97%) [12-44]. PCB could be performed CT, ultrasound or fluoroscopy guidance, in order to target tumor area [14, 18, 20, 21, 26-28, 31, 34-36, 38, 40-44]. In the last years, new types of PCB guided were proposed like MRI-guided biopsy, PET/CT-guided biopsy, or radionuclide-guided biopsy; however, these techniques are costly and are not used routinely in clinical practice [45-47].

\*Correspondence to: Pietro Ruggieri, M.D., Ph.D., Department of Orthopaedics and Orthopaedic Oncology, University of Padova, Via Giustiniani 2, 35128, Padova, Italy; Tel: 390498213372, 393333266234; E-mail: [pietro.ruggieri@unipd.it](mailto:pietro.ruggieri@unipd.it)

The aim of this study was to review our experience with imaging-guided PCB for musculoskeletal lesions in order to define if this could be defined as the gold standard for diagnosis, evaluating 1) percentage of diagnostic procedures, 2) diagnostic accuracy and 3) incidence of complications.

### Case Series

This is a retrospective series of imaging-guided PCB performed at our institution between January 2016 and September 2019. PCB was performed in all cases under imaging guidance: using 14-gauge (1, 62 mm) semi-automatic tru-cut needle guided by ultrasound in soft tissue lesions (322 cases), or with 8-gauge (3, 26 mm) core needle in bone lesions guided by fluoroscopy (376 cases). Sites are summarized in (Tables 1 & 2). For each patient, we reviewed data regarding sex, age, site, history and imaging, expected diagnosis, type of biopsy, complications, histopathology report of biopsy and final histopathology report in those subsequently treated.

**Table 1:** Anatomical distribution of ultrasound-PCB sites in 322 patients with soft tissue lesions.

| Site                      | Number     |
|---------------------------|------------|
| <b>Upper limb</b>         | <b>83</b>  |
| Shoulder                  | 16         |
| Arm                       | 19         |
| Elbow                     | 14         |
| Forearm                   | 15         |
| Wrist                     | 3          |
| Hand                      | 16         |
| <b>Lower limb</b>         | <b>205</b> |
| Hip                       | 16         |
| Thigh                     | 85         |
| Knee                      | 37         |
| Leg                       | 28         |
| Ankle                     | 10         |
| Foot                      | 29         |
| <b>Axial skeleton</b>     | <b>34</b>  |
| Trunk                     | 33         |
| Neck                      | 1          |
| <b>Total biopsy sites</b> | <b>322</b> |

All PCBs were performed after accurate patient history, clinical and radiological evaluations according to 3 main principles. First, a biopsy was located along the surgical approach for future limb-salvage resection in order to be able to remove en-bloc the needle track (potentially contaminated by tumor cells) at the time of definitive resection. Second, the biopsy was performed avoiding contamination of compartments not involved by tumor to prevent cell dissemination that could determine the necessity of major tissue removal during further surgery. Third, areas of the lesion where biopsy was performed were chosen according to preoperative MRI or CT scan, selecting zones with contrast enhancement, avoiding necrotic tissue, including capsule or pseudo-capsule. A mean of 7 samples was performed (range, 6-10), and the material was transferred to our Pathological Institute (and to microbiological evaluation when osteomyelitis is suspected).

**Table 2:** Anatomical distribution of fluoroscopy-PCB sites in 376 patients with bone lesions.

| Site                      | Number     |
|---------------------------|------------|
| <b>Upper limb</b>         | <b>93</b>  |
| Humerus                   | 77         |
| Ulna                      | 8          |
| Radius                    | 5          |
| Phalanges                 | 3          |
| <b>Lower limb</b>         | <b>212</b> |
| Femur                     | 139        |
| Patella                   | 1          |
| Tibia                     | 56         |
| Fibula                    | 6          |
| Calcaneus                 | 3          |
| Metatarsals               | 3          |
| Phalanges                 | 4          |
| <b>Axial skeleton</b>     | <b>71</b>  |
| Scapula                   | 10         |
| Clavicle                  | 5          |
| Acromion                  | 1          |
| Sternum                   | 1          |
| Ribs                      | 1          |
| Vertebrae                 | 1          |
| Pelvis                    | 35         |
| Acetabulum                | 17         |
| <b>Total biopsy sites</b> | <b>376</b> |

Pathologists specialized in orthopaedic oncology evaluated all specimens for the nature of the lesion and specific histological diagnosis. We classified specimens as "diagnostic" (sufficient material) or "non-diagnostic" (insufficient material), according to the histological diagnosis. Moreover, we compared histopathological results from the biopsy with the definitive one in patients subsequently treated surgically. In these cases, we also evaluated Sensitivity, Specificity, Positive Predictive Value (PPV), Negative Predictive Value (NPV), and Diagnostic Accuracy.

In soft tissue, PCB was diagnostic in 304 cases (304/322, 94.4%). In non-diagnostic cases, the biopsy was repeated as PCB in 14 cases and as IB in 4 cases (tumor close to vessels and nerves). In the 14 repeated PCB, diagnostic specimens were obtained in 11, while 3 required further IB. At least 315 PCBs were diagnostic (98%). Diagnoses of soft tissue lesions were reported in (Table 3). One-hundred-nineteen patients were subsequently surgically treated, and in 95% of cases (113/119), there was concordance between histological diagnoses. In 7 cases, final diagnosis differed from biopsy diagnosis: one well-differentiated liposarcoma (vs. soft tissue chondroma), two schwannomas (vs. fibromatosis and pigmented villonodular synovitis), one angiomyxoma (vs. fibromyxoma), one atypical lipomatous tumor (vs. myxofibrosarcoma), one angiolipoma (vs. non-tumor disease) and one infundibular cyst (vs. pigmented villonodular synovitis). PCB in soft tissue has proven to have the ability to distinguish for the nature of the lesions (tumor vs. non-tumor disease) with a Sensitivity of 99%, Specificity of 94%, Diagnostic Accuracy of 98%, PPV of 99% and NPV of 94%. Diagnostic Accuracy for diagnosing benign tumors was 96%, with a Sensitivity of 97%, Specificity of 96%, PPV of 97%, and NPV of 96%. Diagnostic Accuracy

for diagnosing malignant tumors was 99%, with a Sensitivity of 97%, Specificity of 100%, PPV of 100%, and NPV of 98%.

**Table 3:** Final diagnosis of 322 ultrasound-PCB in soft tissue lesions.

| <i>Lesion</i>                                                                     | <i>Number</i> |
|-----------------------------------------------------------------------------------|---------------|
| <b>Benign lesion</b>                                                              | <b>211</b>    |
| Lipoma                                                                            | 68            |
| Hemangioma                                                                        | 31            |
| Pigmented Villo-Nodular Synovitis                                                 | 20            |
| Fibromatosis                                                                      | 18            |
| Schwannoma                                                                        | 14            |
| Cyst                                                                              | 11            |
| Synovial Chondromatosis                                                           | 7             |
| Elastofibroma                                                                     | 5             |
| Others (e.g. Fibromyxoma, Neurofibroma, Fibrolipoma)                              | 37            |
| <b>Malignant lesion</b>                                                           | <b>50</b>     |
| Undifferentiated Pleomorphic Sarcoma                                              | 8             |
| Extraskelatal Chondrosarcoma                                                      | 6             |
| Liposarcoma                                                                       | 6             |
| Metastatic tumor                                                                  | 6             |
| Lymphoma                                                                          | 5             |
| Myxofibrosarcoma                                                                  | 4             |
| Atypical Lipomatous Tumor                                                         | 2             |
| Rhabdomyosarcoma                                                                  | 2             |
| Myxoid liposarcoma                                                                | 2             |
| Synovial Sarcoma                                                                  | 2             |
| Others (e.g. Melanoma, Extraskelatal Osteosarcoma, Extraskelatal Ewing's Sarcoma) | 7             |
| <b>Non-tumor (e.g. Infection, Granuloma, Pseudotumor)</b>                         | <b>43</b>     |
| <b>Undiagnostic</b>                                                               | <b>18</b>     |
| <i>PCB repeated in 14 cases</i>                                                   |               |
| <b>Benign lesion</b>                                                              | <b>6</b>      |
| Hemangioma                                                                        | 3             |
| Lipoma                                                                            | 1             |
| Pigmented Villo-Nodular Synovitis                                                 | 1             |
| Fibromatosis                                                                      | 1             |
| <b>Malignant lesion</b>                                                           | <b>2</b>      |
| Undifferentiated pleomorphic sarcoma                                              | 1             |
| Lymphoma                                                                          | 1             |
| <b>Non-tumor (e.g. Granuloma)</b>                                                 | <b>3</b>      |
| <b>Undiagnostic</b>                                                               | <b>3</b>      |

In bone lesions, PCB was diagnostic in 366 (366/376, 97.3%). In non-diagnostic cases, the biopsy was repeated as PCB in 7 cases and as IB in 3 cases. In the 7 repeated PCBs, diagnostic samples were obtained in 6. At least 372 PCB were diagnostic (99%). Diagnoses of bone lesions were reported in (Table 4). One-hundred-seventy-three patients were subsequently surgically treated, and in 98% of cases (170/173), there was concordance between histological diagnoses. In 3 cases, the final diagnosis differed from biopsy diagnosis: one lymphoma (vs. osteosarcoma), one chondrosarcoma grade 2 (vs. chondroma), and one plasmacytoma (vs. diffuse large B cell lymphoma). PCB in bone lesions has proven to have the ability to distinguish for the nature of the lesions (tumor vs. non-tumor disease) with a Sensitivity of 100%, Specificity of

100%, Diagnostic Accuracy of 100%, PPV of 100% and NPV of 100%. Diagnostic Accuracy for diagnosing benign tumors was 99%, with a Sensitivity of 100%, Specificity of 99%, PPV of 98.5%, and NPV of 100%. Diagnostic Accuracy for diagnosing malignant tumors was 99%, with a Sensitivity of 99%, Specificity of 100%, PPV of 100%, and NPV of 99%. Complications, such as hematoma or wound infection, were not observed after PCB.

**Table 4:** Final diagnosis of 376 fluoroscopy-PCB in bone lesions.

| <i>Lesion</i>                                                                                  | <i>Number</i> |
|------------------------------------------------------------------------------------------------|---------------|
| <b>Benign lesion</b>                                                                           | <b>136</b>    |
| Chondroma                                                                                      | 48            |
| Giant Cell Tumor                                                                               | 17            |
| Aneurysmal Bone Cyst                                                                           | 21            |
| Fibrous dysplasia                                                                              | 14            |
| Chondroblastoma                                                                                | 5             |
| Others (e.g. Histiocytic fibroma, Intraosseous lipoma, Periosteal chondroma)                   | 31            |
| <b>Malignant lesion</b>                                                                        | <b>152</b>    |
| Osteosarcoma                                                                                   | 29            |
| Chondrosarcoma                                                                                 | 20            |
| Metastatic Tumor                                                                               | 68            |
| Lymphoma/ Myeloma                                                                              | 30            |
| Others (e.g. Leiomyosarcoma, Biphasic synovial sarcoma)                                        | 5             |
| <b>Non-tumor (e.g. Bone necrosis, Granuloma, Bone marrow reconversion, Metabolic diseases)</b> | <b>78</b>     |
| <b>Undiagnostic</b>                                                                            | <b>10</b>     |
| <i>PCB repeated in 7 cases</i>                                                                 |               |
| <b>Benign lesion</b>                                                                           | <b>2</b>      |
| Fibrous dysplasia                                                                              | 1             |
| Hemangioma                                                                                     | 1             |
| <b>Malignant lesion</b>                                                                        | <b>1</b>      |
| Ewing's sarcoma                                                                                | 1             |
| <b>Non-tumor (e.g. Bone necrosis)</b>                                                          | <b>3</b>      |
| <b>Undiagnostic</b>                                                                            | <b>1</b>      |

## Discussion

Obtaining a specific histological diagnosis is mandatory before the treatment of musculoskeletal lesions [1-6]. Different types of biopsy could be used with related advantages and disadvantages [3-6, 12-44]. PCB is reported to be a safe procedure burdened by few minor complications only; however, its diagnostic accuracy is lower than with IB due to the limited material obtainable, especially in soft tissue lesions usually presenting necrotic tissue, that makes anatomopathological diagnosis more complicated (Table 5) [12-44]. For this reason, incisional biopsy is still used in some Specialized Centers.

The aim of our study was to review our experience with PCB for the diagnosis of musculoskeletal lesions in order to evaluate if this type of biopsy could be defined as the gold standard. All PCBs were imaging-guided to provide significant/adequate samples containing vital tumor cells targeting the correct tumor area based on preoperative studies (CT or MRI with contrast).

**Table 5:** Summary of the published studies reporting on PCB for diagnosis of musculoskeletal tumors: systematic review.

| <i>Study</i>           | <i>Pt (n)</i> | <i>Bone (n) or Soft tissue (n)</i> | <i>Imaging-guidance</i>                   | <i>Gauge of needle</i> | <i>Passes (n)</i> | <i>Diagnostic results after CNB (%)</i>                                           | <i>Diagnostic accuracy after surgical treatment</i>                                                                                               | <i>Complications</i>                  |
|------------------------|---------------|------------------------------------|-------------------------------------------|------------------------|-------------------|-----------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|
| Zornoza, 1982 [35]     | 42            | Soft tissue                        | CT or US                                  | 18 or 14               | >2                | 83% (35/42)                                                                       | -                                                                                                                                                 | None                                  |
| Skrzynsky, 1996 [14]   | 62            | Bone (17)<br>Soft tissue (45)      | None                                      | -                      | 3-6               | 84% (52/62)                                                                       | DA 100% in Bone<br>DA 78% in Soft tissue                                                                                                          | Track infection (1)                   |
| Schweitzer, 1996 [17]  | 138           | Bone                               | FS                                        | 12 or 18               | 2-3               | 98% (135/138)                                                                     | -                                                                                                                                                 | -                                     |
| Heslin, 1997 [19]      | 60            | Soft tissue                        | None                                      | -                      | -                 | 93% (56/60)                                                                       | Cc in diagnosis malignancy (56/56)<br>Cc in diagnosing grade (40/56)                                                                              | -                                     |
| Dupuy, 1998 [31]       | 176           | Bone<br>Soft tissue                | CT                                        | 14                     | -                 | -                                                                                 | Cc 93% (164/176)                                                                                                                                  | Hematoma (1)<br>Vasovagal symptom (1) |
| Yao, 1999 [33]         | 141           | Bone (56)<br>Soft tissue (85)      | CT (122)<br>FS (13)<br>US (4)<br>None (2) | 12-14<br>14-18         | 3-8               | Bone 75% (42/56)<br>Soft tissue: 82% (70/85)                                      | Cc 73% (41/56) in Bone<br>Cc 75% (64/85) in Soft tissue                                                                                           | None                                  |
| Welker, 2000 [32]      | 161           | Bone (83)<br>Soft tissue (78)      | None (90)<br>CT (55)<br>FS (28)           | 14-17 / 8-11<br>14-15  | 3-5               | 88% (142/161)                                                                     | DA nature of the lesion 92%<br>DA grade 89%<br>DA specific diagnosis 73%                                                                          | Hematoma (1)<br>Drainage (1)          |
| Konermann, 2000 [38]   | 65            | Bone<br>Soft tissue                | US                                        | 14                     | 3-5               | 94% (61/65)                                                                       | -                                                                                                                                                 | None                                  |
| Torriani, 2002 [36]    | 65            | Bone (27)<br>Soft tissue (38)      | US                                        | 14                     | >5                | 96% (63/65)                                                                       | Cc 97% (47/48)                                                                                                                                    | None                                  |
| Jelinek, 2002 [22]     | 110           | Bone                               | CT (85)<br>FS (25)                        | 7-14 / 12-16<br>14-18  | 3-10              | 88% (97/110)                                                                      | Cc 98% (89/91) in diagnosis malignancy<br>Cc 91% (83/91) in diagnosing grade                                                                      | Hematoma (1)                          |
| Issakov, 2003 [40]     | 215           | Bone (135)<br>Soft tissue (80)     | CT                                        | 11-14<br>14-18         | 3-10              | Bone 87% (118/135)<br>Soft tissue 94% (75/80)                                     | -                                                                                                                                                 | Hematoma (3)                          |
| Ray-Coquard, 2003 [30] | 110           | Soft tissue                        | CT or US                                  | 14                     | 4                 | 94% (103/110)                                                                     | Cc 88% (91/103)                                                                                                                                   | Bleeding (1)<br>Hematoma (6)          |
| Yang, 2004 [15]        | 50            | Bone<br>Soft tissue                | -                                         | -                      | 6                 | 98% (49/50)                                                                       | DA nature of the lesion 94% (47/50)<br>DA specific diagnosis 86% (43/50)<br>DA histologic grading 87% (27/31)<br>DA histologic typing 92% (46/50) | None                                  |
| Mitsuyoshi, 2006 [18]  | 163           | Bone (91)<br>Soft tissue (72)      | None (119)<br>CT (44)                     | 16<br>-                | >2                | Bone 88% (80/91)<br>Soft tissue 88% (63/72)<br>No guidance: 85%<br>CT-guided: 93% | Cc 88% (126/143)                                                                                                                                  | Hematoma (1)                          |
| Battaglia, 2007 [44]   | 164           | Soft tissue                        | US                                        | 14-18 or 13-14         | 3-4               | D 90% (148/164)                                                                   | -                                                                                                                                                 | -                                     |
| Sung, 2009 [41]        | 309           | Bone (167)                         | US (151)                                  | -                      | -                 | Bone 92%                                                                          | In 185 surgically treated                                                                                                                         | -                                     |

|                      |      | Soft tissue (142)               | CT (89)<br>FS (69)              |                                |      | Soft tissue 89%                                 | DA 89% in Bone<br>DA 79% in Soft tissue                                                                                                                                   |                                         |
|----------------------|------|---------------------------------|---------------------------------|--------------------------------|------|-------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|
| Kasraeian, 2010 [12] | 57   | Soft tissue                     | None                            | 14                             | 3-5  | 86% (49/57)                                     | Cc 81%                                                                                                                                                                    | None                                    |
| Adams, 2010 [27]     | 233  | Bone (16)<br>Soft tissue (217)  | None                            | -                              | 1-10 | 94% (219/233)                                   | Cc 97% in diagnosing malignancy<br>Cc 81% in diagnosis and grade                                                                                                          | None                                    |
| Strauss, 2010 [37]   | 530  | Soft tissue                     | None                            | -                              | >4   | 93% (493/530)                                   | In 371 surgically treated<br>DA 98% in diagnosing sarcomas<br>DA 86% in diagnosing grade<br>DA 88% in diagnosing benign subtype<br>DA 88% in diagnosing malignant subtype | Hematoma (1)<br>Bleeding (1)            |
| Yang, 2010 [29]      | 508  | Bone (272)<br>Soft tissue (236) | CT (339)<br>US (169)            | 16 or 20                       | 1-6  | Bone 87% (237/272)<br>Soft tissue 96% (226/236) | -                                                                                                                                                                         | -                                       |
| Rimondi, 2011 [28]   | 2027 | Bone                            | CT                              | 8/15                           | >2   | 77% (1567/2027)                                 | -                                                                                                                                                                         | Transient paresis (18)<br>Haematoma (4) |
| Peer, 2011 [42]      | 223  | Soft tissue                     | US                              | 14-18                          | 3-10 | 95% (211/223)                                   | In 113 surgically treated<br>DA 100% in diagnosing malignancy                                                                                                             | -                                       |
| Pohlig, 2012 [13]    | 46   | Bone (33)<br>Soft tissue (13)   | CT or US                        | 14                             | 3-5  | Bone 100% (33/33)<br>Soft tissue 92% (12/13)    | DA 100% in Bone<br>DA 85% in Soft tissue                                                                                                                                  | None                                    |
| Seng, 2013 [43]      | 134  | Bone<br>Soft tissue             | CT                              | -                              | 3-5  | 95% (127/134)                                   | Cc 88% (118/134)<br>DA 94% (31/33) in Bone<br>DA 96% (71/74) in Soft tissue                                                                                               | Wound bruising                          |
| Nouh, 2014 [34]      | 49   | Bone                            | CT                              | 16-18 or 12-15                 | 6    | 88% (43/49)                                     | Cc 100% (29/29)                                                                                                                                                           | None                                    |
| Trieu, 2016 [26]     | 1131 | Bone (380)<br>Soft tissue (751) | CT                              | 14 or 18                       | -    | Bone 88% (334/380)<br>Soft tissue 94% (703/751) | DA 81% in Bone<br>DA 83% in Soft tissue                                                                                                                                   | 0.8%                                    |
| Walker, 2018 [39]    | 105  | Soft tissue                     | None                            | -                              | -    | -                                               | In 69 surgically treated<br>Cc 87%<br>DA 94% nature of the lesion                                                                                                         | Hematoma (2)<br>Bleeding (1)            |
| Current study        | 722  | Bone (400)<br>Soft tissue (322) | FS (376)<br>US (322)<br>CT (24) | 8 in Bone<br>14 in Soft tissue | 6-10 | Bone 97% (366/376)<br>Soft tissue 94% (304/322) | Cc 170/173 (98%) in Bone<br>Cc 113/119 (95%) in Soft tissue                                                                                                               | None                                    |

Pt: Patients; -: Not Reported; PCB: percutaneous biopsy; US: Ultrasound; CT: Computed Tomography; FS: Fluoroscopy; D: Diagnostic; Cc: Concordance between histology of biopsy and of surgery; DA: Diagnostic Accuracy.

This study had some limitations. First, this is a retrospective nonrandomized series, in which we perform PCB for bone as well as soft tissue lesions. However, PCB was performed with a tru-cut needle guided by ultrasound in all soft tissue lesions and with a core needle guided by fluoroscopy in all bone lesions. Second, PCB was performed to detect tumor as well as to exclude it. Consequently, not all patients underwent surgery, according to the treatment required by histological diagnosis, limiting our series and statistical power of analysis. However, non-surgically treated patients were periodically followed in the out-patient clinic, and no cases of misdiagnosis were observed. Third, diagnoses were heterogeneous, precluding evaluation of diagnostic accuracy related to histotypes. However, these kinds of tumors are rare, and we do not preclude further study with more specific analysis.

Imaging-guided PCB can allow diagnosis in most musculoskeletal lesions. Bony PCBs are usually fluoroscopy-guided and performed with a trocar of 11 Gauge, with a percentage of diagnostic results ranging between 75% to 98% [14, 17, 18, 22, 27, 31-33, 41]. Instead, PCBs for soft tissue lesions are generally ultrasound-guided and performed with a 14-gauge tru-cut needle, with a diagnostic result reported in literature ranging from 83% to 96% [12, 13, 19, 20, 29, 32, 33, 35-44]. The incidence of non-diagnostic PCB is higher in soft tissue than in bone lesions, probably due to the high presence of necrotic areas that could make histological interpretation more difficult [34, 43, 44]. In our series, adequate material for diagnosis was obtained more frequently in bone (97%, 366/376), than in soft tissue (94%, 304/322). Repeating another PCB led to diagnosis in 99% and 98% of cases, respectively in bone and soft tissue lesions: non-diagnostic results could be defined as sampling error, due to necrotic, fibrous, or haemorrhagic tissue. In order to reduce non-diagnostic procedures, multiple samples from different areas of the tumor should be done using larger needles [44].

Our results seem to be better than those reported in the literature obtaining only 2.7% and 5.6% of non-diagnostic samples, respectively in bone (we used 8 Gauge trocar) and soft tissue (we performed at least 6 samples). Moreover, targeting with CT the proper area of the lesion to obtain vital tumor cells could be useful. Mitsuyoshi *et al.* [18] compared CT guided and fluoroscopy-guided PCB in 163 patients and report a decrease of non-diagnostic results with CT compared with fluoroscopy (7% vs. 14%), concluding that CT-guidance is more efficacious, thanks to high contrast resolution and spatial definition. However, since CT is more expensive and exposes to a high dose of radiation, its use should be limited.

Also, the diagnostic accuracy is usually higher for bone tumors than for soft-tissue masses, ranging from 80% to 100% and from 76% -100%, respectively [13-15, 18, 26-28, 35-38, 40-42]. The diagnosis of bone lesions can be aided by imaging tools, whereas soft-tissue lesions have more tumor necrosis and more different diagnoses [13]. In our series, diagnostic accuracy was determined by comparing histopathological results from PCB and subsequent tumor resection, in patients that subsequently were surgically treated. We identified an overall diagnostic accordance of 98% in bone lesions, and 95% in soft tissue tumors.

Complications are frequent (up to 17%) when IB is performed [6, 11, 13, 14, 18, 23]. Mankin *et al.*, were the first in 1982 to report the percentages of complications that occurred following an open biopsy in 329 patients

[6]. In 17% of cases, there were hematomas, infections, surgical wound dehiscence, and dispersion of tumor cells in adjoining tissues. In 8.5% of patients, it was found that this procedure negatively affected the prognosis. Finally, it emerged that 4.5% of the patients underwent amputation following the open biopsy [6]. Conversely, complications after PCB are infrequent (less than 2%) for both bone and soft tissue lesions and are usually minor complications, such as biopsy tract infection, post-procedural bleeding, transient paresis [14, 15, 26-28, 35-39]. According to the literature, in our series, we did not encounter any complications related to PCB.

In conclusion, PCB is a safe, minimally invasive, and cost-effective technique for the diagnosis of bone and soft tissue lesions. Indications should be carefully evaluated by an experienced orthopaedic oncologist concerning the suspected entity, size, and location of the lesion to avoid incorrect or deficient results. Targeting the proper area of the lesion to obtain vital tumor cells is mandatory; it is based on pre-biopsy careful imaging evaluation (especially MRI) and imaging-guidance during the procedure. Obtaining multiple samples from different areas of the tumor is essential to have more representative specimens. IB should be reserved for "difficult" cases or after previous non-diagnostic PCBs.

## REFERENCES

1. Potter BK, Adams SC, Pitcher JD Jr, Temple HT (2008) Local Recurrence of Disease After Unplanned Excisions of High-Grade Soft Tissue Sarcomas. *Clin Orthop Relat Res* 466: 3093-3100. [[Crossref](#)]
2. Campanacci M (1999) The Wrong Approach to Tumors of the Musculo Skeletal System: What Should Not Be Done. *Chir Organi Mov* 84: 1-17. [[Crossref](#)]
3. Bickels J, Jelinek JS, Shmookler BM, Neff RS, Malawer MM (1999) Biopsy of Musculoskeletal Tumors. Current Concepts. *Clin Orthop Relat Res* 368: 212-219. [[Crossref](#)]
4. Shives TC (1993) Biopsy of Soft-Tissue Tumors. *Clin Orthop Relat Res* 289: 32-35. [[Crossref](#)]
5. Leithner A, Maurer Ertl W, Windhager R (2009) Biopsy of Bone and Soft Tissue Tumours: Hints and Hazards. *Recent Results Cancer Res* 179: 3-10. [[Crossref](#)]
6. Mankin HJ, Lange TA, Spanier SS (1982) The Hazards of Biopsy in Patients With Malignant Primary Bone and Soft-Tissue Tumors. *J Bone Joint Surg Am* 64: 1121-1127. [[Crossref](#)]
7. Angelini A, Calabrò T, Pala E, Trovarelli G, Maraldi M *et al.* (2015) Resection and Reconstruction of Pelvic Bone Tumors. *Orthopedics* 38: 87-93. [[Crossref](#)]
8. Mavrogenis AF, Pala E, Angelini A, Ferraro A, Ruggieri P (2013) Proximal Tibial Resections and Reconstructions: Clinical Outcome of 225 Patients. *J Surg Oncol* 107: 335-342. [[Crossref](#)]
9. Pala E, Trovarelli G, Calabrò T, Angelini A, Abati CN (2015) Survival of Modern Knee Tumor Megaprotheses: Failures, Functional Results, and a Comparative Statistical Analysis. *Clin Orthop Relat Res* 473: 891-899. [[Crossref](#)]
10. Trovarelli G, Cappellari A, Angelini A, Pala E, Ruggieri P (2019) What Is the Survival and Function of Modular Reverse Total Shoulder Prostheses in Patients Undergoing Tumor Resections in Whom an Innervated Deltoid Muscle Can Be Preserved? *Clin Orthop Relat Res* 477: 2495-2507. [[Crossref](#)]

11. Cannon S, Dyson P (1987) Relationship of the site of open biopsy of malignant bone tumors to local recurrence following resection and prosthetic replacement. *J Bone Joint Surg* 11: 825-832.
12. Kasraeian S, Allison DC, Ahlmann ER, Fedenko AN, Menendez LR (2010) A Comparison of Fine-Needle Aspiration, Core Biopsy, and Surgical Biopsy in the Diagnosis of Extremity Soft Tissue Masses. *Clin Orthop Relat Res* 468: 2992-3002. [[Crossref](#)]
13. Pohlig F, Kirchhoff C, Lenze U, Schauwecker J, Burkart R et al. (2012) Percutaneous Core Needle Biopsy Versus Open Biopsy in Diagnostics of Bone and Soft Tissue Sarcoma: A Retrospective Study. *Eur J Med Res* 17: 29. [[Crossref](#)]
14. Skrzynski MC, Biermann JS, Montag A, Simon MA (1996) Diagnostic Accuracy and Charge-Savings of Outpatient Core Needle Biopsy Compared With Open Biopsy of Musculoskeletal Tumors. *J Bone Joint Surg Am* 78: 644-649. [[Crossref](#)]
15. Yang YJ, Damron TA (2004) Comparison of Needle Core Biopsy and Fine-Needle Aspiration for Diagnostic Accuracy in Musculoskeletal Lesions. *Arch Pathol Lab Med* 128: 759-764. [[Crossref](#)]
16. Barth RJ Jr, Merino MJ, Solomon D, Yang JC, Baker AR (1992) A Prospective Study of the Value of Core Needle Biopsy and Fine Needle Aspiration in the Diagnosis of Soft Tissue Masses. *Surgery* 112: 536-543. [[Crossref](#)]
17. Schweitzer ME, Gannon FH, Deely DM, O'Hara BJ, Juneja V (1996) Percutaneous Skeletal Aspiration and Core Biopsy: Complementary Techniques. *AJR Am J Roentgenol* 166: 415-418. [[Crossref](#)]
18. Mitsuyoshi G, Naito N, Kawai A, Kunisada T, Yoshida A et al. (2006) Accurate Diagnosis of Musculoskeletal Lesions by Core Needle Biopsy. *J Surg Oncol* 94: 21-27. [[Crossref](#)]
19. Heslin MJ, Lewis JJ, Woodruff JM, Brennan MF (1997) Core Needle Biopsy for Diagnosis of Extremity Soft Tissue Sarcoma. *Ann Surg Oncol* 4: 425-431. [[Crossref](#)]
20. Kim SY, Chung HW, Oh TS, Lee JS (2017) Practical Guidelines for Ultrasound-Guided Core Needle Biopsy of Soft-Tissue Lesions: Transformation from Beginner to Specialist. *Korean J Radiol* 18:361-369. [[Crossref](#)]
21. Hau A, Kim I, Kattapuram S, Hornicek FJ, Rosenberg AE et al. (2002) Accuracy of CT-guided Biopsies in 359 Patients With Musculoskeletal Lesions. *Skeletal Radiol* 31: 349-353. [[Crossref](#)]
22. Jelinek JS, Murphey MD, Welker JA, Henshaw RM, Kransdorf MJ et al. (2002) Diagnosis of Primary Bone Tumors With Image-Guided Percutaneous Biopsy: Experience With 110 Tumors. *Radiology* 223: 731-737. [[Crossref](#)]
23. Schwartz HS, Spengler DM (1997) Needle Tract Recurrences After Closed Biopsy for Sarcoma: Three Cases and Review of the Literature. *Ann Surg Oncol* 4: 228-236. [[Crossref](#)]
24. Davies NM, Livesley PJ, Cannon SR (1993) Recurrence of an Osteosarcoma in a Needle Biopsy Tract. *J Bone Joint Surg Br* 75: 977-978. [[Crossref](#)]
25. Saghie S, Masrouha KZ, Musallam KM, Mahfouz R, Abboud M et al. (2010) The Risk of Local Recurrence Along the Core-Needle Biopsy Tract in Patients With Bone Sarcomas. *Iowa Orthop J* 30: 80-83. [[Crossref](#)]
26. Trieu J, Schlicht SM, Choong PF (2016) Diagnosing Musculoskeletal Tumours: How Accurate Is CT-guided Core Needle Biopsy? *Eur J Surg Oncol* 42: 1049-1056. [[Crossref](#)]
27. Adams SC, Potter BK, Pitcher DJ, Temple HT (2010) Office-based Core Needle Biopsy of Bone and Soft Tissue Malignancies: An Accurate Alternative to Open Biopsy With Infrequent Complications. *Clin Orthop Relat Res* 468: 2774-2780. [[Crossref](#)]
28. Rimondi E, Rossi G, Bartalena T, Ciminari R, Alberghini M et al. (2011) Percutaneous CT-guided Biopsy of the Musculoskeletal System: Results of 2027 Cases. *Eur J Radiol* 77: 34-42. [[Crossref](#)]
29. Yang J, Frassica FJ, Fayad L, Clark DP, Weber KL (2010) Analysis of Nondiagnostic Results After Image-Guided Needle Biopsies of Musculoskeletal Lesions. *Clin Orthop Relat Res* 468: 3103-3111. [[Crossref](#)]
30. Ray Coquard I, Ranchère Vince D, Thiesse P, Ghesquière H, Biron P et al. (2003) Evaluation of Core Needle Biopsy as a Substitute to Open Biopsy in the Diagnosis of Soft-Tissue Masses. *Eur J Cancer* 39: 2021-2025. [[Crossref](#)]
31. Dupuy DE, Rosenberg AE, Punyatabandhu T, Tan MH, Mankin HJ (1998) Accuracy of CT-guided Needle Biopsy of Musculoskeletal Neoplasms. *AJR Am J Roentgenol* 171: 759-762. [[Crossref](#)]
32. Welker JA, Henshaw RM, Jelinek J, Shmookler BM, Malawer MM (2000) The Percutaneous Needle Biopsy Is Safe and Recommended in the Diagnosis of Musculoskeletal Masses. *Cancer* 89: 2677-2686. [[Crossref](#)]
33. Yao L, Nelson SD, Seeger LL, Eckardt JJ, Eilber FR (1999) Primary Musculoskeletal Neoplasms: Effectiveness of Core-Needle Biopsy. *Radiology* 212: 682-686. [[Crossref](#)]
34. Nouh MR, Abu Shady HM (2014) Initial CT-guided Needle Biopsy of Extremity Skeletal Lesions: Diagnostic Performance and Experience of a Tertiary Musculoskeletal Center. *Eur J Radiol* 83: 360-365. [[Crossref](#)]
35. Zomoza J, Bernardino ME, Ordóñez NG, Thomas JL, Cohen MA (1982) Percutaneous Needle Biopsy of Soft Tissue Tumors Guided by Ultrasound and Computed Tomography. *Skeletal Radiol* 9: 33-36. [[Crossref](#)]
36. Torriani M, Etchebehere M, Amstalden E (2002) Sonographically Guided Core Needle Biopsy of Bone and Soft Tissue Tumors. *J Ultrasound Med* 21: 275-281. [[Crossref](#)]
37. Strauss DC, Qureshi YA, Hayes AJ, Thway K, Fisher C et al. (2010) The Role of Core Needle Biopsy in the Diagnosis of Suspected Soft Tissue Tumours. *J Surg Oncol* 102: 523-529. [[Crossref](#)]
38. Konermann W, Wuisman P, Ellermann A, Gruber G (2000) Ultrasonographically Guided Needle Biopsy of Benign and Malignant Soft Tissue and Bone Tumors. *J Ultrasound Med* 19: 465-471. [[Crossref](#)]
39. Walker JB, Stockwell E, Worhacz K, Kang P, Decomas A (2018) Safety and Accuracy of Core Needle Biopsy for Soft Tissue Masses in an Ambulatory Setting. *Sarcoma* 2018: 1657864. [[Crossref](#)]
40. Issakov J, Flusser G, Kollender Y, Merimsky O, Lifschitz Mercer B et al. (2003) Computed Tomography-Guided Core Needle Biopsy for Bone and Soft Tissue Tumors. *Isr Med Assoc J* 5: 28-30. [[Crossref](#)]
41. Sung KS, Seo SW, Shon MS (2009) The Diagnostic Value of Needle Biopsy for Musculoskeletal Lesions. *Int Orthop* 33: 1701-1706. [[Crossref](#)]
42. Peer S, Freus T, Loizides A, Gruber H (2011) Ultrasound Guided Core Needle Biopsy of Soft Tissue Tumors; A Fool Proof Technique? *Med Ultrason* 13: 187-194. [[Crossref](#)]
43. Seng C, Png W, Tan MH (2013) Accuracy of Core Needle Biopsy for Musculoskeletal Tumours. *J Orthop Surg (Hong Kong)* 21: 92-95. [[Crossref](#)]

- 
44. Battaglia M, Pollastri P, Ferraro A, Betoni F, Bacci G et al. (2007) The Role of Ultrasound-Guided Needle Biopsy in the Diagnosis of Soft-Tissue Tumors. *J Ultrasound* 10: 59-62. [[Crossref](#)]
  45. Carrino JA, Khurana B, Ready JE, Silverman SG, Winalski CS (2007) Magnetic Resonance Imaging-Guided Percutaneous Biopsy of Musculoskeletal Lesions. *J Bone Joint Surg Am* 89: 2179-2187. [[Crossref](#)]
  46. Win AZ, Aparici CM (2013) Real-time FDG PET/CT-guided bone biopsy in a patient with two primary malignancies. *Eur J Nucl Med Mol Imaging* 40: 1787-1788.
  47. von Meyenfeldt EM, Siebenga J, van der Pol HAG, Schreurs WMJ, Hulsewe KWE (2014) Radionuclide-guided Biopsy of Bone Lesions in Cancer Patients; A Reliable, Well-Tolerated Technique. *Eur J Surg Oncol* 40: 193-196. [[Crossref](#)]