

Available online at www.sciencerepository.org

Science Repository



Research Article

Regulation of Herbal Medicine in Hungary

Gabriella Hegyi^{1*} and Anna Blázovics²

¹Pecs University, Hungary

²Semmelweis University, Budapest, Hungary

ARTICLE INFO

Article history:

Received: 24 January, 2020

Accepted: 18 February, 2020

Published: 28 February, 2020

Keywords:

Medical plants

current methods

human studies

ABSTRACT

In order to meet the growing needs of the European Union medicinal and aromatic plants, Hungary is still one of the major exporters today, although earlier, until the 1980s, we were considered a great herb-growing nation. Therefore, the development of the national medicinal and aromatic herb growing sector is an essential requirement for maintaining and developing our market position. The natural conditions of Hungary favor the cultivation of medicinal and aromatic plants. There are more than 330 herbs in our country. About seventy percent of the drugs grown and harvested are exported. However, significant changes are needed in domestic sales and further processing in order to become a leading industrial sector. Hungary also possesses modern separation techniques and large instruments for analysis, which allow the proper analytical examination of drugs and the precise determination of their active substances. Our universities and research institutes are also prepared to study the active ingredients of herbs by molecular biological methods, which may allow the exact mechanism of action of bioactive substances to be determined and the herbs to be reassessed.

Our accession to the European Union the palette of products made from former herbs has significantly changed. From 2013, the category of medicinal product was discontinued, and the preparations were classified as either food or medicine. However, there are significant costs associated with qualifying as a drug, so drugs or extracts are largely added to dietary supplements. As a result, the strange situation is that dietary supplements are prohibited from making any health claims, even though the dietary supplement is actually not for the benefit of a healthy population, but for those who have a health problem or do not have suitable foods, which are rich in bioactive substances. The justification of phytotherapy must be demonstrated, depending on the state of the art. In the context of safe therapy, studies must follow the requirements of medicines and only then can they become herbal medicines. There is still much to be desired in this area in our country. Classification as a drug is based on expensive animal tests and clinical studies. A new problem arises here, as herbal cure is currently part of the arsenal of complementary medicine. With few exceptions, clinical doctors do not use herbal therapies. Patients usually ask the advice of pharmacists, naturalists, and more rarely, dietitians.

At the Arteriosclerosis Research Group, at the II. Department of Medicine and Institute of Pharmacognosy several decades of biomedical research have served and serve the purpose of evaluating and reassessing the effects of plants used in folk medicine. Over the years, we have been searching for drugs, extracts, and medicines of natural origin that are useful in liver and intestinal diseases known in folk medicine, and are believed to have lipid lowering, antioxidant and immune stimulating effects. In the article, animal and human studies will be also presented.

© 2020 Gabriella Hegyi. Hosting by Science Repository.

*Correspondence to: Dr. Gabriella Hegyi, M.D., Ph.D., Pecs University, Hungary; E-mail: drhegyi@hu.inter.net

Drug Growing and Research in Brief

In order to meet the growing needs of the European Union medicinal and aromatic plants, Hungary is still one of the major exporters today, although earlier, until the 1980s, we were considered a great herb-growing nation. Therefore, the development of the national medicinal and aromatic herb growing sector is an essential requirement for maintaining and developing our market position. The natural conditions of Hungary favor the cultivation of medicinal and aromatic plants. There are more than 330 herbs in our country. Of the cultivated herbs, chamomile, mint and lavender are of paramount importance. The largest areas are mustard, poppy, fennel and coriander, as well as St. John's wort and lemongrass. Many herbs are grown farming on large-scale methods, such as cumin, anise or mustard. Marjoram or marigold can be grown in smaller areas with greater manual labor.

About seventy percent of the drugs grown and harvested are exported. The major buyers are Spain and Germany. However, significant changes are needed in domestic sales and further processing in order to become a leading industrial sector. Hungary also possesses modern separation techniques and large instruments for analysis, which allow the proper analytical examination of drugs and the precise determination of their active substances. Our universities and research institutes are also prepared to study the active ingredients of herbs by molecular biological methods, which may allow the exact mechanism of action of bioactive substances to be determined and the herbs to be reassessed.

The Problem of Becoming Herbal Medicine

Our accession to the European Union the palette of products made from former herbs has significantly changed. From 2013, the category of medicinal product was discontinued, and the preparations were classified as either food or medicine.

However, there are significant costs associated with qualifying as a drug, so drugs or extracts are largely added to dietary supplements. As a result, the strange situation is that dietary supplements are prohibited from making any health claims, even though the dietary supplement is actually not for the benefit of a healthy population, but for those who have a health problem or do not have suitable foods, which are rich in bioactive substances. The justification of phytotherapy must be demonstrated, depending on the state of the art. In the context of safe therapy, studies must follow the requirements of medicines and only then can they become herbal medicines. There is still much to be desired in this area in our country. Classification as a drug is based on expensive animal tests and clinical studies.

A new problem arises here, as herbal cure is currently part of the arsenal of complementary medicine. With few exceptions, clinical doctors do not use herbal therapies. Patients usually ask the advice of pharmacists, naturalists, and more rarely, dietitians. Unfortunately, the worst case scenario is the most common when they are self-medicated on the basis of uncertain sources. In this case either they do not go to the doctor or they stop speaking about the herbal medicine.

It is very difficult to find right way for patients, even the doctors and pharmacists. Today in Hungary, at the beginning of August 2019, there are 17,313 dietary supplements with a notification number, and 497 dietary supplements have been listed as prohibited products by the National Institute for Food and Nutrition Science (OÉTI), one of the General Directorates of National Institute of Pharmacy and Nutrition (OGYÉI).

OÉTI performs professional-methodological, scientific, research, training, professional supervision and expert tasks in the field of nutrition and food. In addition to its statutory tasks, the Institute also performs the task of notification and expertise of food supplements and foods for particular nutritional uses. The Institute's main goal is to achieve healthy nutrition and related prevention in the widest possible range of the Hungarian population.

Medical Biochemical Research at Hungary (Semmelweis University)

At the Arteriosclerosis Research Group, at the II. Department of Medicine and Institute of Pharmacognosy several decades of biomedical research have served and serve the purpose of evaluating and reassessing the effects of plants used in folk medicine. Over the years, we have been searching for drugs, extracts, and medicines of natural origin that are useful in liver and intestinal diseases known in folk medicine, and are believed to have lipid lowering, antioxidant and immune stimulating effects. These agents often have choleric, cholekinetic, and intestinal motility enhancing effects.

We are adapting and developing state-of-the-art, accurate assays for in vitro and in vivo herbal extracts, herbal medicines and dietary supplements. Over the years, we have used spectrophotometric, luminometric, HPLC, GC, GC/MS, HPLC-DAD-MS/MS, OPTLC, ESR, impulse radiolysis and ICP-OES techniques to answer our questions [1-13]. Direct and indirect methods were used to determine the antioxidant and scavenger properties of bioactive molecules. Spectrophotometric and luminometric methods were developed to investigate the antioxidant property in vitro and in vivo. The results obtained with these methods were compared with ESR, pulse radiolysis and amperometric techniques [14]. Luminometric methods suitable for routine examinations have also been developed in humans [15-17]. In our animal experiments, biochemical, immunological and quantitative analytical studies were supplemented with morphological studies [3, 18, 19].

Light and electron microscopy images confirmed tissue damage caused by free radicals and the beneficial physiological effects of natural antioxidants. Preparations were investigated in short term animal experiments (alimentary hyperlipidemia, partial hepatectomy, cannulation of the ductus choledochus, passage-excluded bowel prepare with circulatory) and in human studies lasting several months [15, 16, 20-22].

In our experiments, several plant extracts of natural origin, such as *Raphanus sativus niger*, *Sempervivum tectorum*, *Cichorium intibus*, *Taraxacum officinale*, *Helichrysum arenarium*, *Peroselinum crispum*, *Anthriscus cerefolium* lyophilized extracts, *Solidago canadensis* and

alcoholic and aqueous extracts of *Ginkgo biloba*, *Hibiscus sabdariffa*, *Beta vulgaris var rubra* as well as silibinin and silymarin compounds or fruits, apples sour cherries were investigated [7, 12, 18, 19, 23-41].

Among the products, vitamin E, Hegrimarin, Raphacol, Rowachol and Detralex were tested [15, 16, 20, 22, 39, 42-46]. We performed a partial phytochemical and elemental analysis of Beiqishen tea, Holub, Rose and Mate teas, as well as examined several tea with special function (Tisztító/Cleansing, Salaktalanító/ Decontaminating, Gyomor-tea/Stomach Tea) and green tea products were analyzed [11, 47-49]. These studies confirmed our hypothesis that pharmaceutical preparations are often so-called "drug bombs". In many cases, product advertising is templated and does not describe any side effects that may occur.

Early we observed changes the effects of free radical reactions in the different liver diseases. In addition to fatty liver and extrahepatic cholestasis models, we also introduced the partial hepatectomy model in our research repertoire [3, 20, 40]. In animal experiments, we confirmed the importance of redox homeostasis and the importance of free radical - antioxidant balance [13, 50]. In our research we studied the free radical changes in bile in connection with stone formation and the irritant effect of abnormal bile on the intestinal mucosa and their prevention with natural active substances [8, 19, 22]. Histopathological examinations were applied to study the morphological changes [51, 52]. These studies have brought us closer to a better understanding of the intestinal-liver axis [53]. To investigate the effect of plant-based active substances and metal ions on their free radical antioxidant status, a research repertoire was developed. During the research several drugs, lyophilisates, tea and medicines were examined [15, 16, 20, 25, 38, 44, 47, 49, 54-56].

Elemental metabolism is altered in hyperlipidemia. Most toxic elements are accumulated, indicating a decrease in the liver's detoxifying capacity. Therefore, natural active substances that are capable of accelerating the depletion of these metal ions may serve as tools for primary prevention [8, 9, 54]. Polyphenols can remove heavy metal ions from the body [57]. Recent studies have confirmed that the beneficial physiological effect of *Sempervivum tectorum* also involves the elimination of toxic metal ions accumulated in the liver [54, 8].

The therapeutic use of multicomponent natural products or their enjoyment in the form of enjoyment raises several problems. I emphasize that plants can accumulate different metal ions, but it is not excluded that toxic elements are introduced into the preparations due to soil contamination [9, 48, 58, 59]. It is also unfortunate that the refreshing effect is often provided by increasing the caffeine content, so that some preparations may contain unreasonably high levels of caffeine [6, 49].

This brief summary focuses on the mechanism of action of the lyophilised aqueous extract of *Sempervivum tectorum* in rat experiments, vacuum dried beetroot, dietary supplement containing vegetable and fruit concentrate and Raphacol bile granulate in human studies to demonstrate the broad applied methodological approach.

Sempervivum Tectorum

The question was how can the polyphenol type antioxidants of *Sempervivum tectorum* act in hyperlipidemia and in fatty liver? The

lyophilized aqueous extract of *Sempervivum tectorum* contains significant amounts of polyphenolic compounds resulting in superoxide anion and hydrogen peroxide / hydroxyl radical scavenger properties. This extract protects the bowel epithelial cells against damaging effects of lipids (lipid peroxides) in hyperlipidemia. The cellular immune functions changed considerably during partial hepatectomy. *Sempervivum tectorum* modifies cytokine production in fatty liver in partial hepatectomy. We observed that this lyophilized extract had a favorable effect on concanavalin-A-stimulated spleen cell blast transformation, lipopolysaccharide-triggered TNF- α production and spontaneous IL-1 activity in partial hepatectomy.

Conclusion: *Sempervivum tectorum* has several beneficial effects on alimentary induced fatty liver and on fatty liver after partial hepatectomy. The consumption of plant-based foods has a beneficial effect on the intestinal redox balance [19, 23].

Vacuum Dried *Beta vulgaris L. ssp. esculenta var. rubra*

Beta vulgaris L. ssp. esculenta var. rubra is rich in bioactive compounds therefore it affects numerous biochemical reactions, enzyme activities and metabolic pathways, therefore this vegetable can be considered as functional food.

Question: How does betaine content of *Beta vulgaris L. ssp. esculenta var. rubra* diminish the anemia? Among the quaternary ammonium compounds, which have human health significance due to their biological activity, a high concentration of betaine can be found in the table beet.

Betaine is an important natural molecule for treating homocystinuria, alcoholic steatosis, chemically induced liver, lung and skin cancers. This molecule helps to create choline, can help synthesis of carnitine and helps to convert homocysteine into methionine, and it takes part in biologic methylation.

Methylation is very important for hemoglobin synthesis. Betaine is a potential HCHO generator, therefore serves the HCHO to the methylation of globin to stabilize hemoglobin formation, considering the arginine (38), methionine (65) and the lysine (72) near the methionine (80) must be methylated and coordinated towards the central iron of heme. Folic acid plays a catalytic role in transmethylation.

In this clinical study 10 g natural table beet vacuum dried product was given twice daily for 1 month for 24 patients (mean age 68 \pm 8 years) with hormone-resistant and metastatic prostate cancer treated with taxane chemotherapy, who reported their complaints themselves first, mean 3.6 \pm 2.8 years before. 18 men's data were amenable after treatment for evaluation.

In addition to routine laboratory examination values of HbA1c, 9 cytokines and levels of 3 growth factors, the global parameters of redox-homeostasis, few elements, Zn- and free protoporphyrin levels, trans-methylation processes were determined before and one month after treatment. The measured parameters were improved during treatment. Zn- and free protoporphyrin levels were decreased and transmethylation ability was better as well as redox parameters, but PSA and EGF

concentrations were higher more than 50% of patients in the end of the treatment.

Conclusion: Results clearly verify that iron, folic acid and betaine components as well as colorful compounds with antioxidant activity of table beet extract demand more attention as a preventive therapy in chemotherapy induced anemia. Table beet will have a great impact and application in human cancer, but because of the increasing values of EGF and PSA close medical control is necessary for patients especially during chemotherapy [55].

Fruit and Vegetable Color Concentration

In this study the question was: How can a dietary supplement act on the colectomised patients in secondary prevention?

We studied the effect of antioxidant and metal element rich proprietary preparation "Fruit and vegetable color concentration"- which contained bioactive agents - on colectomised volunteers. Components of „Fruit and vegetables color concentration” are: *Sambucus nigra* (berry), *Vaccinium oxycoccus* (extract), *Morus nigra* (berry), *Beta vulgaris var. rubra*, *Vitis vinifera* (berry), *Hippophae rhamnoides* (fruit oil, berry juice), *Aronia rotundifolia* (extract), *Hibiscus sabdariffa* (extract). Beneficial medical effects are due to bioactive components, such as tansmethylation agent betaine, betanins, betaxanthins, vulgaxanthine, polyphenols, flavonoids, vitamins (thiamine (B1), riboflavine (B2), piridoxine (B6), ascorbic acid, biotin and folic acid) as well as soluble fiber, pectin and different metal elements (e.g. Al, B, Ba, Ca, Cu, Fe, K, Mg, Mn, Na, Zn), which act on various physiological routes. According to newer researches, these colored pigments have antioxidant activity and play an important role in the development of antioxidant status in the human organism.

This concentration (2x3 g/day) was added to 28 adult patients as secondary prevention after 5-10 years of their colectomy and 9 healthy volunteers for 3 months. Several routine laboratory parameters were not altered after treatment compared to start time data in colectomised patients and control volunteers. The treatment did not change the tumor marker levels significantly, but in each case the values were elevated in examined groups. Redox parameters of colectomised patients showed strengthened free radical reactions, namely rebound effect against antioxidant excess, although tendency was observed in controls as well. Anti-inflammatory effects of the natural product were not clearly observed. The dietary supplement treatment negative influenced the plasma and erythrocyte free radical levels and the HbA1c concentration, diminished the reducing power similarly reduced the GSHPx- level in the asymptomatic patients. The HbA1c levels well correlated with free radical levels of erythrocytes and the lowered antioxidants status.

Tumor marker AFP, CEA and CA-19-9 levels of operated patients were higher in most cases, and only in few cases the normal values exceeded comparison with baseline ones [60, 61].

Conclusion: Moderate nutritional customs with natural scavengers or antioxidants can help to restore the normal function of tissues and organs, but the immoderate consumption of vitamins and other bioactive agents, such as polyphenols, flavonoids, isothiocyanates etc. are contraindicated [62].

Raphacol

Hypothesis: The anti-inflammatory, lipid peroxidation lowering and radical scavenging properties as well as gallstone dissolving effect of black radish root may be beneficial in reducing of bowel discomfort of patients with IBD. The active ingredients of the preparation can significantly influence the redox state of intestinal epithelial cells. The mild bactericidal, fungicidal action can protect the normal intestinal flora of the intestine. The question was, while what time does the beneficial effect of treatment occur and does the unwanted side effect occur during several months of treatment? Patients were under regular medical control and received adequate treatment of IBD. Blood samples were collected before the first administration of Raphacol (0.2 g/day), and after 3, 6, 9 and 12 months of treatment.

No significant differences were observed in serum parameters during the 12 months of Raphacol treatment in the moderately severe condition of IBD during the term. At the end of the treatment, IL-6 level was decreased, which was also measurable at the end of 3 months after treatment, and free SH-group concentrations increased. Redox-parameters were deteriorated in the plasma. Undesirable changes of sera bile acid were detected by the end of the period. Raphacol significantly reduced erythrocyte chemiluminescence and moderately reduced erythrocyte HbA1c levels by month 9, which was favorable. From the data obtained, it can be concluded that the Raphacol granule composition can exert a long-term influence on the redox homeostasis even at 0.2 g/day concentration. It probably modifies redox reactions at several levels and at several points. New redox homeostasis was formed.

Conclusion: Continued use of Raphacol is not recommended [63].

Short Discussion

The presented studies are very important for the safe therapeutic uses of natural products. In Hungarian universities in this field the examinations are continued in several diseases, especially in tumors with drug extracts and isolated bioactive agents, considering the several failure of targeted cancer therapy

Conflicts of Interest

None.

Abbreviations

EGF: Epidermal growth factor

ESR: Electron spin resonance spectroscopy

GC: Gas chromatography

GC/MS: Gas chromatography–mass spectrometry

HPLC: High-performance liquid chromatography

HPLC-DAD-MS/MS: High-performance liquid chromatography with diode-array detection tandem mass spectrometry

ICP-OES: Inductively coupled plasma optical emission spectroscopy

OPTLC: Over pressured thin-layer chromatography

PSA: Prostate specific antigen

REFERENCES

1. György I, Antus S, Blázovics A, Földiák G (1992) Substituent effects in the free radical reactions of silybin radiation-induced oxidation of the flavonoid at neutral pH. *Int J Radiat Biol* 61: 603-609. [Crossref]
2. György I, Blázovics A, Fehér J, Földiák G (1990) Reactions of inorganic free radicals with liver protecting drugs. *Radiat Phys Chem* 36: 165-167.
3. Blázovics A, Fehér E, Fehér J (1992) Role of free radical reactions in experimental hyperlipidemia in the pathomechanism of fatty liver. *Free Radicals and Liver* 98-126.
4. Blázovics A, Prónai L, Kéry Á, Petri G, J Fehér (1993) Natural antioxidant extract from *Sempervivum tectorum*. *Phytother Res* 7: 95-97.
5. Blázovics A, Lugasi A, Kemény T, Hagymási K, Kéry Á (2000) Membrane stabilizing effects of natural polyphenols and flavonoids from *Sempervivum tectorum* on hepatic microsomal mixed-function oxidase system in hyperlipidemic rats. *J Ethnopharm* 73: 479-485. [Crossref]
6. Blázovics A, Lugasi A, Szentmihályi K, Kéry Á (2003) Reducing power of the natural polyphenols of *Sempervivum tectorum* in vitro and in vivo. *Acta Biol Szegediensis* 47 4: 99-102.
7. Lugasi A, Dworschák E, Blázovics A, Kéry Á (1998) Antioxidant and free radical scavenging properties of squeezed juice from black radish (*Raphanus sativus* L. Var, niger) root. *Phytotherapy Res* 12: 502-506.
8. Szentmihályi K, Blázovics A, Lugasi A, Kéry Á, Lakatos B et al. (2000) Effect of natural polyphenol-type antioxidants (*Sempervivum tectorum* and *Raphanus sativus* L. var. niger extracts) on metal ion concentrations in rat bile fluid. *Current Topics in Biophysics* 24: 203-207.
9. Szentmihályi K, Fehér E, Vinkler P, Kéry Á, Blázovics A (2004) Metabolic alterations of toxic and none essential elements by the treatment of *Sempervivum tectorum* extract in hyperlipidemic rat model. *Toxicol Pathol* 32: 50-57.
10. Héthelyi É, Szarka Sz, Antal I, Blázovics A, Galambosi B et al. (2004) *Gentiana lutea* L. fitokémiai- és biokémiai vizsgálata. Olaj, Szappan, Kozmetika, (Phytochemical and biochemical examination of *Gentiana lutea* L., Oil, Soap, Cosmetics) 53: 204-211.
11. Héthelyi É, Szabó L, Rapavi E, Domokos J, Blázovics A (2001) Tényeken alapuló orvoslás a fitoterápiában. Olaj, Szappan, Kozmetika, (Evidence-based medicine in phytotherapy, Oil, Soap, Cosmetics) 50: 67-75.
12. Balázs A, Tóth M, Blazics B, Héthelyi É, Szarka Sz et al. (2012) Investigation of dietary important components in selected red fleshed apples by GC-MS and LC-MS. *Fitoterapia* 83: 1356-1363. [Crossref]
13. Blázovics A, Sárdi É (2018) Methodological repertoire development to study the effect of dietary supplementation in cancer therapy. *Microchem J* 136: 121-127.
14. Prónai L, Blázovics A, Horváth ÉM, Láng I, Fehér J (1993) Superoxide activity of dihydroquinoline type derivative (CH 402 and MTDQ-DA). *Free Rad Res Comms* 19: 287-296. [Crossref]
15. Lugasi A (2004) Élelmiszer eredetű antioxidánsok hatása primer és szekunder prevencióban: Állatkísérletes és humán tanulmányok. Ph.D. Tézisek, Budapest, 2004. (The effect of food-derived antioxidants in primary and secondary prevention: Animal and human studies. Ph.D. Theses, Budapest, 2001.)
16. Kocsis I (2004) A redox-homeosztázis vizsgálati lehetőségei, jelentősége állatkísérletes és humán tanulmányokban. Ph.D. Tézisek, Budapest, 2004. (Examination possibilities of redox homeostasis and significance in animal and human studies. Ph.D. Theses, Budapest, 2004.)
17. Hagymási K, Kocsis I, Lugasi A, Fehér J, Blázovics A (2002) Extrahepatic biliary obstruction: can sylimarin protect liver function? *Phytother Res* 1: S78-S80. [Crossref]
18. Blázovics A, Fehér E, Kéry Á, Petry G, Fehér J (1993) Liver protecting and lipid lowering effects of *Sempervivum tectorum* in the rats. *Phytother Res* 7: 98-100.
19. Blázovics A, González-Cabello R, Barta I, Gergely P, Fehér J (1994) Effect of liver-protecting *Sempervivum tectorum* extract on the immune reactivity of spleen cells in hyperlipidemic rats. *Phytother Res* 8: 33-37.
20. Horváth ÉM, Gonzalez-Cabello R, Blázovics A, van der Looij M, Barta I et al. (2001) Effect of silibinin and vitamin E on restoration of cellular immune response after partial hepatectomy. *J Ethnopharmacol* 77: 227-232. [Crossref]
21. Lugasi A, Blázovics A, Lebovics VK, Kocsis I, Kéry Á (2002) Beneficial health effect of black radish and in vivo experimental conditions. *Phytochemistry and Pharmacology* 2: 365-375.
22. Sípos P (2001) Az epe szabadgyökös változásai epekövességben és hatása a bélnyálkahártyára. Ph.D. Tézisek. (Free radical changes of bile in gallstone diseases and their effect on intestinal mucosa. Ph.D. Theses, 2001.)
23. Blázovics A, Horváth ÉM, Barta I, Kéry Á, Petri G et al. (1993) *Sempervivum tectorum* and liver regeneration, In: Proc. of the Oxygen free radicals and scavengers in the natural sciences. Akadémiai Kiadó, Budapest 293-300.
24. Kocsis I, Hagymási K, Kéry Á, Szőke É, Blázovics A (2003) Effect of chicory on pancreas status of rats in experimental hyperlipidemia. *Acta Biol Szegediensis* 47: 143-146.
25. Kocsis I, Szentmihályi K, Rapavi E, Hubay M, Kéry Á et al. (2004) Effect of *Cichorium intibus* (L.) on the redox balance and metal ion content of fatty liver in hyperlipidemic rats. *Trace Elements and Electrolytes* 21: 262-268.
26. Hagymási K, Blázovics A, Lugasi A, Kristó SzT, Fehér J et al. (2000) In vitro antioxidant evaluation of dandelion (*Taraxacum officinale* Web) water extract. *Acta Aliment Hung* 29: 1-7.
27. Czinner E, Hagymási K, Blázovics A, Kéry Á, Szőke É et al. (2000) In vitro antioxidant properties of *Helichrysum arenarium* L. Moench. *J Ethnopharmacol* 73: 437-443. [Crossref]
28. Czinner E, Hagymási K, Blázovics A, Kéry Á, Szőke É et al. (2001) The in vitro effect of *Helichrysi flos* on microsomal lipid peroxidation. *J Ethnopharmacol* 77: 31-35. [Crossref]
29. Czinner E, Lemberkovics É, Kéry Á, Hagymási K, Blázovics A et al. (1999) Biologically active components of *Helichrysum arenarium* L. *Eur J Drug Metab Pharmacokinetics* 24: 309-313. [Crossref]
30. Fejes Sz, Kéry Á, Blázovics A, Lugasi A, Lemberkovics É et al. (1998) *Petroselinum crispum* (Mill.) Nym. ex A.W. Hill. in vitro antioxidáns hatásának vizsgálata. *Acta Pharmaceut Hung* 68: 150-156. [Crossref]
31. Fejes Sz, Blázovics A, Lugasi A, Lemberkovics É, Petri G et al. (2000) Free radical scavenging and membrane protecting effect of methanol extracted fraction of Parsley. *Acta Aliment Hung* 29: 81-87.
32. Fejes Sz, Blázovics A, Lemberkovics É, Petri G, Szőke É et al. (2000) Free radical scavenging and membrane protective effects of methanol

- extracts from *Anthriscus cerefolium* and *Petroselinum crispum* (Mill.) Nym. ex. A.W. Hill. *Phytother Res* 14: 362-365. [Crossref]
33. Fejes Sz, Blázovics A, Lugasi A, Lemberkovich É, Petri G et al. (2000) In vitro antioxidant activity of *Anthriscus cerefolium* L. (Hoffm.). *J Ethnopharmacol* 69: 259-265. [Crossref]
 34. Apáti P, Balázs A, Blázovics A, Stevenson G, Houghton PJ et al. (2004) In vitro módszerek alkalmazása *Solidago canadensis* L. lipidperoxidáció gátló hatásának vizsgálatában. A szabadgyök-kutatás aktuális kérdései, MSZKT Munkaértekezlet, (Use of in vitro methods to investigate the inhibitory effect of *Solidago canadensis* L. on lipid peroxidation. Current issues in free radical research, MSZKT Workshop) Budapest.
 35. Rapavi E, González-Cabello R, Szentmihályi K, Székely E, Blázovics A (2006) The effect of calyx infusion of *Hibiscus sabdariffa* L. on T-cells-mediated immune response in mitogen-induced blastogenesis of human lymphocytes in vitro. *Acta Alim Hung* 35: 281-288.
 36. Váli L, Fébel H, Stefanovits-Bányai É, Sárdi É, Lugasi A et al. (2006) Duodenum protecting effects of table beet (*Beta vulgaris* var. *rubra*) during hepatic ischaemia-reperfusion. *Acta Alimentaria* 35: 445-453.
 37. Váli L, Stefanovits Bányai É, Szentmihályi K, Fébel H, Sárdi É et al. (2007) Liver protecting effects of table beet (*Beta vulgaris* var. *rubra*) during ischemia-reperfusion. *Nutrition* 23: 172-178. [Crossref]
 38. Blázovics A, Somogyi A, Nagy É, Fehér J (1988) Silibinin kezelés hatása a máj drog-metabolizáló enzimrendszerére kísérletes hiperlipidaemiában. Kísérletes Orvostudomány, (Effect of silibinin treatment on the drug metabolizing enzyme system of the liver in experimental hyperlipidemia.) *Exp Med* 151-158.
 39. Fehér J, Blázovics A, González-Cabello R, Horváth M É, Gregely P (1996) Effects of silibinin and vitamin E on lipid peroxidation and cellular immunoreactivity in experimentally induced fatty liver of rats. *Med Sci Monit* 2: 397-403.
 40. Hagymási K, Blázovics A, Lengyel G, Kocsis I, Fehér J (2001) Oxidative damage in alcoholic liver disease. *Eur J Gastroent Hepatol* 13: 49-53.
 41. Papp N, Blázovics A, Fébel H, Salido S, Altarejos J et al. (2015) Antihyperlipidemic effects of sour cherries characterized by different in vitro antioxidant power and polyphenolic composition. *Plant Foods Human Nutr* 70: 408-413. [Crossref]
 42. Gonzalez-Cabello R, Blázovics A, Horváth ÉM, Müzes Gy, Gergely P et al. (1992-1993) Effect of vitamin E on the immunoreactivity of spleen cells in hyperlipidemic rats. *Acta Med Hung* 49: 101-108. [Crossref]
 43. Ágoston M, Örsi F, Fehér E, Hagymási K, Orosz Zs et al. (2003) Sylimarin and vitamin E reduce amiodarone-induced lysosomal phospholipidosis in rats. *Toxicology* 190: 231-241. [Crossref]
 44. Hagymási K (2002) A máj redox-homeosztázisának tanulmányozása kísérletes és humán vizsgálatokban: antioxidáns kezelés – antioxidáns betegség. PhD-tézisek. Budapest. (Study of liver redox homeostasis in experimental and human studies: antioxidant treatment - antioxidant disease. PhD Thesis. Budapest.
 45. Sipos P, Hagymási K, Lugasi A, Fehér E, Blázovics A (2002) Effect of black radish root (*Raphanus sativus* L. var *niger*) on the colon mucosa in rats fed a fat rich diet. *Phytotherapy Res* 16: 677-679. [Crossref]
 46. Rapavi E, Szentmihályi K, Kocsis I, Stefanovits-Bányai É, Blázovics A (2002) The effect of hesperidin and diosmin on ion concentration and antioxidant defense system in fatty liver in rats. Metal elements in environment, medicine and biology. 5th International Symposium, November 4-6, 2002, Timisoara, Romania .
 47. Blázovics A, Rapavi E, Hagymási K, Szentmihályi K, Lugasi A et al. (2004) Is Medical control of herbal tea consumption necessary? *Acta Horticulturae* No. 629, ISHS, 153-160.
 48. Szentmihályi K, Ladó K, Rapavi E, Taba G, Kocsis I et al. (2004) Gyógynövények ásványi elemtartalmának vizsgálata és értékelése a komplex prevenciók kutatásokkal összefüggésben. Széchenyi Szimpózium, (Investigation and evaluation of mineral content of medicinal plants in the context of complex prevention research. Széchenyi Symposium) 1/016. NKFP.
 49. Rapavi E, Stefanovits-Bányai É, Szentmihályi K, Lugasi A, Szecsó V, Sztetanov A et al. (2003) Trace elements and antioxidant properties of Hungarian dried herbs. New results in the trace element research. ed.: I. Pais. Szent István University, Budapest 210-220.
 50. Blázovics A, Kovács Á, Lugasi A (2015) The effect of short and long term antioxidant treatments on redox homeostasis in experimental and clinical studies. in: V. Watkins Sarah (Ed.), Nova Science Publisher, 2007, 2015, pp. 1-34 Chapter 4.
 51. Fehér E, Blázovics A, Horváth ÉM, Kéry Á, Petry G, Fehér J (1993) Effect of *Sempervivum tectorum* extract in the experimental hyperlipidemia and alcoholism. A histological study in the liver and jejunum. in: Role of Free Radicals in Biological Systems, ed.: J. Fehér, A. Blázovics, B. Matkovich, M. Mézes, Akadémiai Kiadó, Budapest 45-57.
 52. Blázovics A, Kéry Á, Fehér E, Prónai L, González-Cabello R et al. (1997) Natural antioxidants in liver therapy. *Current Topics Biophysics* 20: 14-30.
 53. Egresi A, Kovács Á, Szilvás Á, Blázovics A (2017) Bél-máj tengely vizsgálata colitis ulcerosaban – retrospektív tanulmány. (Bowel-liver axis examination in ulcerative colitis - retrospective study). *Orv Heti* 158: 1014-1021. [Crossref]
 54. Blázovics A, Lugasi A, Hagymási K, Szentmihályi K, Kéry A (2002) Natural antioxidants and tissue regeneration. Curative effect and reaction mechanism. *Phytochem Pharmacol* 2002 8: 93-134.
 55. Blázovics A, Nyirády P, Romics L, Szűcs M, Horváth A et al. (2012) How can cancer-associated anemia be moderated with nutritional factors and how do *Beta vulgaris* L. ssp. *esculenta* var. *rubra* modify the transmethylation reaction in erythrocytes in cancerous patients? in: D.S. Silverberg (Ed.), Anemia, INTECH, 2012, 1–23 Open Access.
 56. Kocsis I, Lugasi A, Hagymási K, Kéry A, Fehér J et al. (2002) Beneficial properties of black radish root (*Raphanus sativus* L. var. *niger*) squeezed juice in hyperlipidemic rats, biochemical and chemiluminescence measurements. *Acta Alim Hung* 32: 185-190.
 57. Blázovics A, Szentmihályi K, Lugasi A, Balázs A, Hagymási K et al. (2003) In vitro analysis of the properties of Beiqishen tea. *Nutrition* 19: 869-875. [Crossref]
 58. Stefanovits-Bányai É, Bertényi-Divinyi Zs, Kiss AS, Blázovics A, Koczka N (2002) Comparison of calcium, magnesium, potassium and sodium contents of *Ginkgo biloba* L. leaves from Hungary. *J Elementol* 7: 49-56.
 59. Stefanovits-Bányai É, Koczka N, Szecsó V, Sztetanov A, Bertényi-Divinyi Zs et al. (2002) Phytochemical and element analysis of *Ginkgo biloba* L. from Hungary. In Proc 10th International race Element Symposium, Budapest, ed.: I. Pais.
 60. Szilvás Á, Blázovics A, Székely Gy, Dinya E, Fehér J et al. (2001) Comparative study between the free radicals and tumor markers in

-
- patients with gastrointestinal tumors. *J Physiology Paris* 95: 247-252. [[Crossref](#)]
61. Szilvás Á, Székely Gy, Tarján Z, Fomet B (2002) Three-dimensional ultrasosography, virtual colonoscopy and endorectal magnetic resonance imaging in the diagnosis of complicated inflammatory bowel disease. *Endoscopy* 34: 94. [[Crossref](#)]
62. Blázovics A, Kursinszki L, Papp N, Kleiner D, Szőke É et al. (2016) Is professional prescription of a commercially derived dietary supplement in colectomysed patients necessary? *Eur J Integr. Med* 8: 219-226.
63. Blázovics A, Kovács Á, Lugasi A (2007) The effect of long term antioxidant treatments on redox homeostasis in clinical studies. *Hungarian Med J* 1: 83-95.