Available online at www.sciencerepository.org

Science Repository



Research Article

Metabolomics of Rat Brain After Treatment with Phenelzine: High-Resolution Mass Spectrometric Demonstration of Increased Brain Levels of N-Acetyl Amino Acids

Paul L. Wood^{1*}, John E. Cebak² and Glen B. Baker³

¹Metabolomics Unit, College of Veterinary Medicine, Lincoln Memorial University, Tennessee, USA

²DeBusk College of Osteopathic Medicine, Lincoln Memorial University, Tennessee, USA

³Neurochemical Research Unit, Department of Psychiatry, University of Alberta, Edmonton, Alberta, Canada

ARTICLE INFO

Article history:

Received: 20 June, 2020

Accepted: 7 July, 2020

Published: 24 July, 2020

Keywords:

Phenelzine

rat brain

N-acetyl amino acids

ABSTRACT

Background: Phenelzine (PLZ) is a non-specific monoamine oxidase inhibitor that has demonstrated clinical efficacy in patients with treatment resistant depression. The mechanism of action with regard to this efficacy is complicated in that its metabolite, β -phenylethylenedihydrazine (PEH), is an inhibitor of amino acid transaminases resulting in dramatic brain elevations of GABA, alanine, ornithine and tyrosine. The full neurochemical profile of PLZ and PEH remain to be explored.

Objective: To undertake a non-targeted metabolomics study of phenelzine on rat brain neurochemistry.

Methods: We undertook a high-resolution mass spectrometric metabolomics analysis of rat cortical brain 1 and 12 hours after intraperitoneal dosing with PLZ or PEH. Tandem mass spectrometry was utilized to obtain relative quantitation data.

Results: N-acetyl amino acids were found to be elevated in cortical brain tissue following either PLZ or PEH treatments.

Conclusions: Our data indicate PLZ treatment significantly augments brain levels of N-acetyl amino acids and that this may involve inhibition of deacylases by PEH and/or induction of N-amino acid acetyltransferases.

© 2020 Paul L. Wood. Hosting by Science Repository.

Introduction

Phenelzine is a nonspecific monoamine oxidase (MAO) inhibitor that has been utilized in patients with treatment resistant depression [1-4]. Phenelzine is a potent inhibitor of both MAO-A and MAO-B which increases brain monoamine levels. In contrast, its metabolite, β -phenylethylenedihydrazine (PEH), is a weak MAO inhibitor but is a potent transaminase inhibitor that significantly elevates brain levels of GABA and alanine, tyrosine, and ornithine [5-8]. PLZ and PEH also are both carbonyl scavenging agents that reduce reactive aldehyde toxicity *in vivo* and *in vitro*. The aldehyde scavenging actions of PLZ and PEH have been hypothesized to be responsible for their neuroprotective actions in animal models of ischemia-reperfusion injury, traumatic brain injury, spinal cord injury, experimental autoimmune encephalitis, and *in vitro* aldehyde toxicity [9-14]. These data indicate that PLZ and PEH

have very complex pharmacodynamic effects. To further investigate the biochemical effects of these compounds, we undertook a non-targeted metabolomics study of PLZ and PEH on the rat brain metabolome. This study revealed that PLZ and PEH significantly augment brain levels of free N-acetyl amino acids.

Methods

I Rat Brain Samples

Male Sprague Dawley rats were dosed intraperitoneally with 30 mg/kg of PLZ or PH, and the brains harvested after decapitation at 1 and 12 hours. The brains were frozen immediately in isopentane on solid carbon dioxide and then removed to containers stored at -80°C until the frontal cortex was dissected out for metabolomics analysis. All procedures

*Correspondence to: Paul L. Wood, Metabolomics Unit, College of Veterinary Medicine, Lincoln Memorial University, 6965 Cumberland Gap Pkwy., Harrogate, Tennessee 37752, USA; E-mail: paul.wood@lmunet.edu

involving animals were approved by the University of Alberta Biosciences Animal Care and Use Committee (AUP00000216) and were in accordance with the guidelines of the Canadian Council on Animal Care.

II Sample Processing

40 to 60 mg of cortical tissue were sonicated in 1 mL of ice-cold acetonitrile:methanol:formic acid (800:200:2.5) containing the stable isotope internal standards [$^2\text{H}_3$]N-acetyl-methionine, [$^2\text{H}_5$]N-acetyl-glutamate, and bromocriptine [15]. After centrifugation at 30,000 x g and 4°C for 30 min, 750 μL of the clear supernatant was dried by centrifugal vacuum evaporation. The samples were next dissolved in acetonitrile:methanol (1:1) for flow infusion analyses.

Table 1: MS² analyses of N-acetyl amino acids.

N-Acetyl Amino Acid	[M-H]	MS ² Product	Product Anion
N-Acetyl Glycine	116.0352	Gly	74.0247
N-Acetyl Proline	156.0665	Pro	114.0717
N-Acetyl Valine	158.0822	Val	116.0717
N-Acetyl Threonine	160.0615	Thr	118.0509
N-Acetyl Hydroxyproline	172.0615	Hydroxyproline	130.0509
N-Acetyl Leucine	172.0979	Leucine	130.0873
N-Acetyl Glutamine	187.0724	Glutamine	145.0618
N-Acetyl Glutamate	188.0564	Glutamate-CO ₂	102.0560
N-Acetyl [$^2\text{H}_5$]Glutamate	193.0731	[$^2\text{H}_5$]Glutamate – CO ₂	107.0873
N-Acetyl-Methionine	190.0547	Methionine	148.0437
N-Acetyl [$^2\text{H}_3$]Methionine	193.0731	[$^2\text{H}_3$]Methionine	151.0626

IV Data Presentation

Data are presented as relative (R) N-acetyl amino acid levels (i.e. the ratio of the signal intensity of the endogenous N-acetyl amino acid to the signal intensity of an appropriate stable isotope internal standard), corrected for protein, \pm SD (N=5).

Results

The most marked observation from our preliminary non-targeted metabolomics analysis was increased brain levels of N-acetyl amino

III High-Resolution Mass Spectrometric Analyses

Samples underwent flow infusion analyses at a flow rate of 12 μL per min. and were analysed via high-resolution mass spectrometry (HR-MS) utilizing a Q-Exactive benchtop orbitrap (Thermo Fisher) with a resolution of 140,000 and less than 1 ppm mass error. Negative ion electrospray ionization (NESI) with a sheath gas of 12, a spray voltage of 3.7 kV, and a capillary temperature of 321°C was used. For the pilot metabolomics analysis, the scan was from 60 to 900 amu. The data were analysed via an in-house Excel (Microsoft) spreadsheet with over 1200 metabolites of interest.

To obtain relative quantitative data of N-acetyl amino acids, MS² studies utilized a window of 0.4 amu for the precursor ion and the product ions were acquired at high resolution (< 1 ppm mass error). For MS² studies (Table 1) the neutral collision energy (NCE) was optimized between 20 and 30 eV.

acids with PLZ and PEH dosing. Since this was a flow infusion analysis, there were a number of potential isobars with the exact mass of each N-acetyl amino acid. Therefore, to obtain relative quantitation data we performed MS² analyses. This approach clearly demonstrated increased levels of an array of brain N-acetyl amino acids (Figure 1). It is interesting to note that N-acetylaspartate levels, which are in millimolar (mM) concentrations in the brain, were unaffected by the drug treatments.




Figure 1: Relative levels (R) of N-acetyl amino acids in rat cortical brain tissue at 1 and 12 hours after treatment with phenelzine (PLZ, 30 mg/kg, i.p.) or β -phenylethylidenehydrazine (PEH). Gly: glycine; Pro: proline; Val: valine; Thr: threonine; Hyp: hydroxyproline; Leu: leucine/isoleucine; Gln: glutamine; Glu: glutamate; Met: methionine. Mean \pm SEM (N=5); *, $p < 0.05$.

Discussion

While our knowledge base regarding post-translational processing of proteins, via N-acetylation of serine, alanine, glycine, methionine, threonine, valine, and aspartate (EC 2.3.1.254-258), or lysine (EC 2.3.1.32) has grown significantly, our understanding of the roles of free N-acetyl amino acids is much more limited [16, 17]. Protein bound N-acetyl amino acids are released by acylaminoacyl peptidase (EC 3.4.19.1); however, free N-acetyl amino acids are also synthesized by a number of N-acetyl transferases. The most studied free N-acetyl amino acid is N-acetylaspartate, since it is present in millimolar concentrations in the brain [18, 19]. N-Acetylaspartate is synthesized by a specific acetylase NAT8L (EC 2.3.1.17) [20, 21]. Whereas N-acetyl glutamate and N-acetylmethionine are both synthesized via amino acid N-acetyltransferase (EC 2.3.1.1) [22, 23]. Interestingly, methamphetamine, a monoamine releaser, has been shown to induce NAT8, the aforementioned synthetic enzyme involved in the biosynthesis of N-acetylaspartate [24, 25]. PLZ, which augments monoamine levels, may therefore also induce NAT8L and augment N-acetyl amino acids. However, PEH which is a weak MAO inhibitor, could not be acting via this mechanism.

Alternatively, inhibition of catabolism might be involved in the augmentation of N-acetyl amino acids, similar to the augmentation of amino acids via inhibition of transaminases by PEH [7]. The potential enzyme targets include aliphatic aminoacylase (EC 3.5.1.14; ACY1) [26, 27], aspartoacylase (EC 3.5.1.15; ACY2), and N-amino aromatic amino acid amidohydrolase (EC 3.5.1.114; ACY3) [28, 29]. In this regard, inhibition of ACY3 provides neuroprotection from aldehyde toxicity, suggesting that in addition to carbonyl scavenging, augmentation of N-acetyl amino acids may contribute to the neuroprotection provided by PLZ in models of neural trauma [9-14, 29].

The functional roles of free N-acetyl amino acids remain to be more fully elucidated but decreased plasma levels of N-acetylmethionine have been monitored in cystic fibrosis patients and childhood obesity, and decreased plasma levels of N-acetyl glycine in obesity [30-33]. Cadmium, which is an ACY1 inhibitor, elevates urinary levels of N-acetylglutamate, N-acetylglutamine, and N-acetylphenylalanine, suggesting a rapid turnover rate for these N-acetyl amino acids *in vivo* [34, 35]. Similarly, precursor labeling studies have defined the rapid dynamics of N-acetylmethionine synthesis in human oligodendrocyte cultures [22].

A detailed analysis of N-acetyl amino acids in dogs with gallbladder mucocele formation found decreased blood levels of N-acetylated alanine, glycine, glutamate, isoleucine, leucine, methionine, serine, and threonine [35]. In contrast, the bile in these dogs was characterized by increased levels of N-acetylated glutamate, histidine, isoleucine, leucine, lysine, threonine, tryptophan, tyrosine, and valine. These data suggest that N-acetyl amino acids play a complex metabolic function in the gallbladder.

With regard to brain function, free N-acetylmethionine, N-acetylglutamine, N-acetylglutamate, N-acetylasparagine, and N-acetylalanine have all been monitored in the human brain [23, 36-40]. N-acetylglutamate is a critical modulator of the urea cycle, while N-acetylleucine modulates the neuronal activity of vestibulocerebellar and

posterolateral thalamic circuits involved in vestibular function, while N-acetylglutamine appears to be involved in the sleep-wake cycle [40-42]. Clearly, we currently have limited knowledge of the roles of N-acetyl amino acids in brain function.

Conflicts of Interest

None.

Funding

This research was funded by Lincoln Memorial University, the University of Alberta, and a grant to GBB from the Canadian Institutes of Health Research (CIHR; grant MOP-86712).

REFERENCES

- Jonathan M Meyer, Michael A Cummings, George Proctor (2017) Augmentation of phenelzine with aripiprazole and quetiapine in a treatment-resistant patient with psychotic unipolar depression: case report and literature review. *CNS Spectr* 22: 391-396. [[Crossref](#)]
- Shulman KI, Herrmann N, Walker SE (2013) Irreversible monoamine oxidase inhibitor (MAOI) antidepressants have significant efficacy in treatment-resistant unipolar depression, Current place of monoamine oxidase inhibitors in the treatment of depression. *CNS Drugs* 27: 789-797.
- Samantha J Thomas, Mirae Shin, Melvin G McInnis, Jolene R Bostwick (2015) Combination therapy with monoamine oxidase inhibitors and other antidepressants or stimulants: strategies for the management of treatment-resistant depression. *Pharmacotherapy* 35: 433-449. [[Crossref](#)]
- David Menkes, Peter Bosanac, David Castle (2016) MAOIs - does the evidence warrant their resurrection? *Australas Psychiatry* 24: 371-373. [[Crossref](#)]
- Kathryn G Todd, Glen B Baker (2008) Neurochemical effects of the monoamine oxidase inhibitor phenelzine on brain GABA and alanine: A comparison with vigabatrin. *J Pharm Pharm Sci* 11: 14s-21s. [[Crossref](#)]
- Dmitriy Matveychuk, Emerson Nunes, Nasir Ullah, Carlos A Velázquez-Martínez, Erin M MacKenzie et al. (2013) Comparison of phenelzine and geometric isomers of its active metabolite, β -phenylethylidenehydrazine, on rat brain levels of amino acids, biogenic amine neurotransmitters and methylamine. *J Neural Transm* 120: 987-996. [[Crossref](#)]
- Dmitriy Matveychuk, Emerson Nunes, Nasir Ullah, Fahad S Aldawsari, Carlos A Velázquez Martínez et al. (2014) Elevation of rat brain tyrosine levels by phenelzine is mediated by its active metabolite β -phenylethylidenehydrazine. *Prog Neuropsychopharmacol Biol Psychiatry* 53: 67-73. [[Crossref](#)]
- Erin M MacKenzie, Suzanne L Grant, Glen B Baker, Paul L Wood (2008) Phenelzine causes an increase in brain ornithine that is prevented by prior monoamine oxidase inhibition. *Neurochem Res* 33: 430-436. [[Crossref](#)]
- Paul L Wood, M Amin Khan, Joseph R Moskal, Kathryn G Todd, Véronique A M I Tanay et al. (2006) Aldehyde load in ischemia-reperfusion brain injury: neuroprotection by neutralization of reactive aldehydes with phenelzine. *Brain Res* 1122: 184-190. [[Crossref](#)]

10. Indrapal N Singh, Lesley K Gilmer, Darren M Miller, John E Cebak, Juan A Wang et al. (2013) Phenzelzine mitochondrial functional preservation and neuroprotection after traumatic brain injury related to scavenging of the lipid peroxidation-derived aldehyde 4-hydroxy-2-nonenal. *J Cereb Blood Flow Metab* 33: 593-599. [[Crossref](#)]
11. Zhe Chen, Jonghyuck Park, Breanne Butler, Glen Acosta, Sasha Vega Alvarez et al. (2016) Mitigation of sensory and motor deficits by acrolein scavenger phenelzine in a rat model of spinal cord contusive injury. *J Neurochem* 138: 328-338. [[Crossref](#)]
12. Curtis A Benson, Grace Wong, Gustavo Tenorio, Glen B Baker, Bradley J Kerr (2013) The MAO inhibitor phenelzine can improve functional outcomes in mice with established clinical signs in experimental autoimmune encephalomyelitis (EAE). *Behav Brain Res* 252: 302-311. [[Crossref](#)]
13. Mee-Sook Song, Glen B Baker, Serdar M Dursun, Kathryn G Todd (2010) The antidepressant phenelzine protects neurons and astrocytes against formaldehyde-induced toxicity. *J Neurochem* 114: 1405-1413. [[Crossref](#)]
14. Glen Baker, Dmitriy Matveychuk, Erin M MacKenzie, Andrew Holt, Yanlin Wang (2019) Attenuation of the effects of oxidative stress by the MAO-inhibiting antidepressant and carbonyl scavenger phenelzine. *Chem Biol Interact* 304: 139-147. [[Crossref](#)]
15. Drazic A, Myklebust LM, Ree R, Arnesen T. (2016) The world of protein acetylation. *Biochim Biophys Acta* 1864: 1372-1401. [[Crossref](#)]
16. Petra Van Damme, Marta Lasa, Bogdan Polevoda, Cristina Gazquez, Alberto Elosegui-Artola et al. (2012) N-terminal acetylome analyses and functional insights of the N-terminal acetyltransferase NatB. *Proc Natl Acad Sci U S A* 109: 12449-12454. [[Crossref](#)]
17. Linda Chang, Joseph Friedman, Thomas Ernst, Kai Zhong, Nicholas D Tsopelas et al. (2007) Brain metabolite abnormalities in the white matter of elderly schizophrenic subjects: implication for glial dysfunction. *Biol Psychiatry* 62: 1396-1404. [[Crossref](#)]
18. Lucia Margari, Andrea De Giacomo, Francesco Craig, Roberto Palumbi, Antonia Pescechera et al. (2018) Frontal lobe metabolic alterations in autism spectrum disorder: a ¹H-magnetic resonance spectroscopy study. *Neuropsychiatr Dis Treat* 14: 1871-1876. [[Crossref](#)]
19. Ariane R Pessentheiner, Helmut J Pelzmann, Evelyn Walenta, Martina Schweiger, Lukas N Groschner et al. (2013) NAT8L (N-acetyltransferase 8-like) accelerates lipid turnover and increases energy expenditure in brown adipocytes. *J Biol Chem* 288: 36040-36051. [[Crossref](#)]
20. Yoshiaki Miyamoto, Noriyuki Iegaki, Kequan Fu, Yudai Ishikawa, Kazuyuki Sumi et al. (2017) Striatal N-acetylaspartate synthetase Shati/Nat8L regulates depression-like behaviors via mGluR3-mediated serotonergic suppression in mice. *Int J Neuropsychopharmacol* 20: 1027-1035. [[Crossref](#)]
21. Ljubica Caldovic, Nicholas Ah Mew, Dashuang Shi, Hiroki Morizono, Marc Yudkoff et al. (2010) N-acetylglutamate synthase: structure, function and defects. *Mol Genet Metab* 100: S13-S19. [[Crossref](#)]
22. Tara Smith, M Said Ghandour, Paul L Wood (2011) Detection of N-acetyl methionine in human and murine brain and neuronal and glial derived cell lines. *J Neurochem* 118: 187-194. [[Crossref](#)]
23. Prasanth S Ariyannur, John R Moffett, Pachiappan Manickam, Nagarajan Pattabiraman, Peethambaran Arun et al. (2010) Methamphetamine-induced neuronal protein NAT8L is the NAA biosynthetic enzyme: implications for specialized acetyl coenzyme A metabolism in the CNS. *Brain Res* 1335: 1-13. [[Crossref](#)]
24. Kyosuke Uno, Toh Miyazaki, Kengo Sodeyama, Yoshiaki Miyamoto, Atsumi Nitta (2017) Methamphetamine induces Shati/Nat8L expression in the mouse nucleus accumbens via CREB- and dopamine D1 receptor-dependent mechanism. *PLoS One* 12: e0174196. [[Crossref](#)]
25. H Lindner, S Höpfner, M Täfler Naumann, M Miko, L Konrad et al. (2000) The distribution of aminoacylase I among mammalian species and localization of the enzyme in porcine kidney. *Biochimie* 82: 129-137. [[Crossref](#)]
26. Jörn Oliver Sass, Jathana Vaithilingam, Corinne Gemperle Britschgi, Cathérine C S Delnooz, Leo A J Kluijtmans et al. (2016) Expanding the phenotype in aminoacylase 1 (ACY1) deficiency: characterization of the molecular defect in a 63-year-old woman with generalized dystonia. *Metab Brain Dis* 31: 587-592. [[Crossref](#)]
27. Jeremy S Francis, Ana Olariu, Scott W McPhee, Paola Leone (2006) Novel role for aspartoacylase in regulation of BDNF and timing of postnatal oligodendrogenesis. *J Neurosci Res* 84: 151-169. [[Crossref](#)]
28. Kirill Tsurulnikov, Natalia Abuladze, Anatol Bragin, Kym Faull, Duilio Cascio et al. (2012) Inhibition of aminoacylase 3 protects rat brain cortex neuronal cells from the toxicity of 4-hydroxy-2-nonenal mercapturate and 4-hydroxy-2-nonenal. *Toxicol Appl Pharmacol* 263: 303-314. [[Crossref](#)]
29. Theresa A Laguna, Cavan S Reilly, Cynthia B Williams, Cole Welchlin, Chris H Wendt (2015) Metabolomics analysis identifies novel plasma biomarkers of cystic fibrosis pulmonary exacerbation. *Pediatr Pulmonol* 50: 869-877. [[Crossref](#)]
30. E Isganaitis, S L Rifas Shiman, E Oken, J M Dreyfuss, W Gall et al. (2015) Associations of cord blood metabolites with early childhood obesity risk. *Int J Obes* 39: 1041-1048. [[Crossref](#)]
31. Steven C Moore, Charles E Matthews, Joshua N Sampson, Rachael Z Stolzenberg Solomon, Wei Zheng (2014) Human metabolic correlates of body mass index. *Metabolomics* 10: 259-269. [[Crossref](#)]
32. Nancy F Butte, Yan Liu, Issa F Zakeri, Robert P Mohny, Nitesh Mehta et al. (2015) Global metabolomic profiling targeting childhood obesity in the Hispanic population. *Am J Clin Nutr* 102: 256-267. [[Crossref](#)]
33. Peter Curley, Chris van der Does, Arnold J M Driessen, Jan Kok, D van Sinderen (2003) Purification and characterisation of a lactococcal aminoacylase. *Arch Microbiol* 179: 402-408. [[Crossref](#)]
34. Sailendra Nath Sarma, Ammar Saleem, Jin Yong Lee, Maki Tokumoto, Gi Wook Hwang et al. (2018) Effects of long-term cadmium exposure on urinary metabolite profiles in mice. *J Toxicol Sci* 43: 89-100. [[Crossref](#)]
35. Jody L Gookin, Kyle G Mathews, John Cullen, Gabriela Seiler (2018) Qualitative metabolomics profiling of serum and bile from dogs with gallbladder mucocele formation. *PLoS One* 13: e0191076. [[Crossref](#)]
36. J V Auditore, L Wade (1973) Isolation of N-acetylglutamine, pyroglutamic acid and the butyl ester of pyroglutamic acid from human brain. *J Neurochem* 21: 335-343. [[Crossref](#)]
37. J V Auditore, L Wade, E J Olson (1966) Occurrence of N-acetyl-L-glutamic acid in the human brain. *J Neurochem* 13: 1149-1155. [[Crossref](#)]
38. J V Auditore, L H Wade (1972) N-acetyl-L-asparagine in human brain. *Neuropharmacology* 11: 385-394. [[Crossref](#)]
39. J V Auditore, L Wade (1971) Occurrence of N-acetylalanine in the human brain. *J Neurochem* 18: 2389-2395. [[Crossref](#)]

-
40. Lisa Günther, Roswitha Beck, Guoming Xiong, Heidrun Potschka, Klaus Jahn et al. (2015) N-acetyl-L-leucine accelerates vestibular compensation after unilateral labyrinthectomy by action in the cerebellum and thalamus. *PLoS One* 10: e0120891. [[Crossref](#)]
 41. Allen K Bourdon, Giovanna Maria Spano, William Marshall, Michele Bellesi, Giulio Tononi et al. (2018) Metabolomic analysis of mouse prefrontal cortex reveals upregulated analytes during wakefulness compared to sleep. *Sci Rep* 8: 11225. [[Crossref](#)]
 42. Wood PL (2020) Flow injection ESI high-resolution mass spectrometry metabolomics analytical platform. IN Springer Protocols, *Neuromethods: Metabolomics* in press (PL Wood, Editor).