



Available online at www.sciencerepository.org

Science Repository



Research Article

Treatment Strategy for Chronic Obstructive Parotitis Related to Diabetes

Chuan-Bin Wu¹ and Lei Xue^{2*}

¹Attending Doctor, Department of Oral and Maxillofacial Surgery, School and Hospital of Stomatology, China Medical University, Liaoning Provincial Key Laboratory of Oral Diseases, PR China

²Department of Oral and Maxillofacial Surgery, School and Hospital of Stomatology, China Medical University, Liaoning Provincial Key Laboratory of Oral Diseases, PR China

ARTICLE INFO

Article history:

Received: 30 March, 2022

Accepted: 15 April, 2022

Published: 3 May, 2022

Keywords:

Diabetes

COP

Sialendoscope

VAS

effect

ABSTRACT

Objectives: The aim of this study was to describe our experience in treating chronic obstructive parotitis related to diabetes.

Methods: Twelve patients with chronic obstructive parotitis (COP) related to diabetes were selected for the study. A sialendoscope was introduced from the orifice to investigate the ductal wall and lumen. During the operation, chymotrypsin and gentamicin were injected. All patients were followed up for 6 months. Preoperative and postoperative visual analog scale (VAS) evaluations and salivary gland scintigraphy (SGS) examinations were applied to evaluate the therapeutic effect; differences were considered statistically significant at $P < 0.05$.

Results: A sialendoscope was successfully used under local anaesthesia in all members of the cohort. As shown by the endoscope, mucus plugs were the most common feature. Some adhered tightly to the ductal wall. We also found ductal congestion in some cases. The postoperative VAS scores and SGS counts were both significantly lower than the preoperative values ($P < 0.05$).

Conclusion: Chymotrypsin administration during interventional sialendoscopy is significantly effective in the treatment of diabetes-related COP.

© 2022 Lei Xue. Hosting by Science Repository.

Get access to the full version of this article: <http://dx.doi.org/10.31487/j.JSO.2022.02.01>

*Correspondence to: Dr. Lei Xue, M.D., Ph.D., Department of Oral and Maxillofacial Surgery, School and Hospital of Stomatology, China Medical University, Liaoning Provincial Key Laboratory of Oral Diseases, Shenyang 110002, Liaoning Province, PR China; Tel: +8602431927735; E-mail: cbwlx8080@163.com