

Available online at www.sciencerepository.org

Science Repository



Case Report

Celecoxib Combined with Thalidomide in the Treatment of Refractory Neoplastic Fever in Advanced Cholangiocarcinoma: A Case Report

Qiu-yuan Liang¹, Kai-Jian Lei^{1,2*}, Shan-Bing Wang², Yue Ren² and Yan Xu²

¹Department of Oncology, The Second People's Hospital of Yibin, Yibin, Sichuan, China

ARTICLE INFO

Article history:

Received: 2 December, 2020 Accepted: 18 December, 2020 Published: 23 December, 2020

Keywords: Celecoxib thalidomide neoplastic fever cholangiocarcinoma

ABSTRACT

Neoplastic fever, a common symptom of tumors, can be mainly controlled by nonsteroidal anti-inflammatories (NSAIDs) and corticosteroids. However, there is no standard treatment guideline for refractory neoplastic fever, which cannot be controlled by the medicines mentioned above. This report presents a case of advanced cholangiocarcinoma with persistent fever lasting 3 weeks. After ineffective treatment with various antibiotics, ibuprofen, and corticosteroids, the body temperature returned to normal after a 48-hour treatment with celecoxib + thalidomide. Finally, the overall survival (OS) of the patient reached 8 months with radiotherapy and chemotherapy.

© 2020 Kai-Jian Lei. Hosting by Science Repository.

Get access to the full version of this article: http://dx.doi.org/10.31487/j.JSO.2020.06.06

²North Sichuan Medical College, Nanchong, Sichuan, China

^{*}Correspondence to: Kai-Jian Lei, Department of Oncology, The Second People's Hospital of Yibin, No. 268, Nanguang Road, Yibin; North Sichuan Medical College, Nanchong, Sichuan, 644000, China; E-mail: lkj600@163.com

^{© 2020} Kai-Jian Lei. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Hosting by Science Repository. http://dx.doi.org/10.31487/j.JSO.2020.06.06