

Available online at www.sciencerepository.org

**Science Repository** 



## Research Article Mechanisms Underlying the Therapeutic Effects of Huangqi, Gegen, Renshen and Sangye in Treating Diabetic Cardiomyopathy Based on Data Mining, Network Pharmacology and Molecular Docking

Xing-Chen Guo<sup>1,3#</sup>, Wan-Hao Gao<sup>1,3#</sup>, Zhi-Wen Zhang<sup>1,3</sup>, Dong-Dong Zhang<sup>2\*</sup> and Mu-Wei Li<sup>1,3\*</sup>

<sup>1</sup>Department of Cardiology, People's Hospital of Zhengzhou University, Zhengzhou University, Zhengzhou, PR China

<sup>2</sup>School of Life Sciences, Shihezi University, Xiangyang Street, Shihezi, PR China

<sup>3</sup>Heart Center of Henan Provincial People's Hospital, No.1 Fuwai Road, Zhengdong New District, Zhengzhou, Henan, 450018, PR China

#Contributed equally

## ARTICLEINFO

Article history: Received: 27 October, 2022 Accepted: 10 November, 2022 Published: 29 November, 2022 Keywords:

Molecular dynamics simulation PI3K-Akt signaling pathway mTOR signaling pathway diabetic cardiomyopathy Puerarin

## ABSTRACT

**Objective:** To evaluate the therapeutic effects of traditional Chinese medicines Radix astragali (Huangqi, HQ), Ginseng (Renshen, RS), Radix puerariae (Gegen, GG), and Mulberry leaf (Sangye, SY) on diabetic cardiomyopathy (DC) based on bioinformatics and network pharmacology, through gene expression analysis of geo clinical samples, molecular docking of compounds and targets, and molecular dynamics simulation, and to discover new targets for prevention or treatment of DC, in order to facilitate and better serve the discovery of new drugs as well as their application in the clinic.

**Materials and Methods:** For the initial selection of ingredients and targets using the TCMSP as a starting point, we performed a primary screening of ingredients and targets of the four herbs using tools including Cytoscape, Tbtools, R 4.0.2, Autodock Vina, PyMOL, and GROMACS. To further screen the effective ingredients and targets, we performed protein interaction network (PPI) analysis (gene = 12), gene expression analysis (n = 24) by clinical samples of DCs from the gse26887 dataset, biological process (BP) analysis (FDR  $\leq$  0.05, gene = 7), KEGG pathway analysis (FDR  $\leq$  0.05, gene = 7), and ingredient target pathway network analysis (gene = 1) by applying these targets from the screen, Biological processes, disease pathways regulated by targets and the relationship between each component target and pathway were obtained. We further screened the targets and visualized the docking results by precision molecular docking of ingredients and targets, after which we performed molecular dynamics simulation and consulted a large number of relevant literature for validation of the results.

**Results:** Through screening, analysis and validation of the data, we finally confirmed the presence of 36 active ingredients in HQ, RS, GG, and SY, which mainly act on AKT1, ADRB2, GSK3B, PPARG, and BCL2 targets, and these five targets mainly regulate PI3K-Akt, Adrenergic signaling in cardiomyocytes, AGE-RAGE signaling pathway in diabetic complications, JAK-STAT, cGMP-PKG, AMPK, and mTOR signaling pathway exert preventive or therapeutic effects on DCM. Molecular dynamics (MD) simulations revealed that the complex formed by Calycosin, Frutinone A, Puerarin, Inophyllum E, the four active components of HQ, RS, GG, and SY, and the four target proteins ADRB2, PPARG, AKT1, and GSK3B acting on DCS is able to exist in a very stable tertiary structure under human environment.

**Conclusion:** Our study successfully explains the effective mechanism of HQ, RS, GG, and SY in ameliorating DC, while predicting the potential targets and active components of HQ, RS, GG, and SY in treating DC, which provides a new basis for investigating novel mechanisms of action at the network pharmacology level and a great support for subsequent DC research.

© 2022 Dong-Dong Zhang & Mu-Wei Li. Hosting by Science Repository.

Get access to the full version of this article: http://dx.doi.org/10.31487/j.JICOA.2022.04.01

<sup>\*</sup>Correspondence to: Dong-Dong Zhang, School of Life Sciences, Shihezi University, Xiangyang Street, Shihezi 832003, PR China; E-mail: sedate@stu.shzu.edu.cn Mu-Wei Li, Department of Cardiology, People's Hospital of Zhengzhou University, Zhengzhou University; Heart Center of Henan Provincial People's Hospital, No.1 Fuwai Road, Zhengdong New District, Zhengzhou, Henan, 450018, PR China; E-mail: lmw0207zzu@163.com

<sup>© 2022</sup> Dong-Dong Zhang & Mu-Wei Li. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Hosting by Science Repository. http://dx.doi.org/10.31487/j.JICOA.2022.04.01