

Available online at www.sciencerepository.org

Science Repository



Review Article

Zebrafish Heart as a Model to Elucidate the Mechanisms of Sudden Cardiac Death

Rafael Antonio Vargas^{1,2*}

ARTICLE INFO

Article history:

Received: 30 November, 2020 Accepted: 21 December, 2020 Published: 31 December, 2020

Keywords:
Zebrafish
sudden cardiac death
channelopathies
cardiomyopathies
development
arrhythmias

ABSTRACT

The heart of zebrafish has been used as a simple, low-cost model to study development, structure and function of the heart from early to late stages of life. Also, it has been established as a model to study different cardiac pathologies generated through different methods. Cardiac pathologies include from functional disorders such as arrhythmia, to structural disorders such as hypertrophic heart disease; in many of them, genetic and molecular aspects have been associated. Noteworthy, some of these genetic and molecular factors have been invoked as causes of sudden cardiac death in humans. This adverse outcome is common in many cardiovascular pathologies and can occur at any age. Unfortunately, etiology and physiopathology of sudden cardiac death are unclear, so extensive research in this area is required. This mini-review highlight three related points: First, the relevance of sudden cardiac death in humans. Second, advances in knowledge of the development, function and pathologies in the zebrafish heart model. Finally, the possibility of using the zebrafish model for the study of sudden cardiac. For this review, a literature search was performed using the PubMed database and the search engine Google Scholar: the words sudden cardiac death, zebrafish, arrhythmias, cardiomyopathies were combined for this search.

© 2020 Rafael Antonio Vargas. Hosting by Science Repository.

Introduction

Sudden cardiac death is the most common lethal outcome of many cardiovascular diseases, and it can occur in any age group, race or sex. The etiology and pathophysiological mechanisms are unknown; however, an important genetic component has been suggested to be involved in the early stages of life, and cardiovascular degenerative changes are decisive in the late stages. This review shows the advantages of zebrafish model considering the genetic, molecular and physiological information available on cardiovascular development and function throughout its life cycle to study sudden cardiac death. Much of the genetic, molecular, physiological and pathophysiological evidence described in zebrafish coincides with the data collected about the possible causes and mechanisms of sudden cardiac death in other animal models and humans.

Literature Review

Extensive literature has been published related to sudden cardiac death, animal models, heart, zebrafish, arrythmias, cardiomyopathies and heart molecular biology (Figure 1). Yet, works contain little or no information on zebrafish as a model to study sudden cardiac death (Figure 2). For this review, the words sudden cardiac death, zebrafish, arrhythmias, cardiomyopathies, channelopathies and heart molecular biology were combined. The number of publications on the topic of sudden cardiac death has been increasing in the last decade (Figures 1 & 2). However, the number of works with zebrafish as a model to study sudden cardiac death is limited; so, this model and this topic have a lot of opportunities for researchers.

¹Facultad de Medicina, Universidad Militar Nueva Granada, Bogotá, Colombia

²Maestria en Actividad Física para la Salud, Universidad Santo Tomás, Bogotá, Colombia

^{*}Correspondence to: Rafael Antonio Vargas, Facultad de Medicina, Universidad Militar Nueva Granada, Maestria en Actividad Física para la Salud, Universidad Santo Tomás, Bogotá, Colombia; Tel: +5713177749194; ORCID: 0000-0001-5702-9240; E-mail: rafael.vargasv@unimilitar.edu.co; rvargas3200@hotmail.com

^{© 2020} Rafael Antonio Vargas. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Hosting by Science Repository. http://dx.doi.org/10.31487/j.JICOA.2020.06.10




Figure 1: Results of literature review conducted from the PubMed database between the years 2000 and 2020. Search was performed with keywords "Rodents and heart", "zebrafish and heart", "rodents and sudden cardiac death" and "zebrafish and sudden cardiac death".




Figure 2: Results of literature review conducted from the PubMed database. Search was performed with keywords "zebrafish and heart molecular biology" (1); "zebrafish and cardiac arrhythmias" (2); "zebrafish and cardiomyopathies" (3); "zebrafish and sudden cardiac death" (4) and "zebrafish and channelopathies" (5). Black bars show total documents and white bars show documents from 2000 to 2020.

Cardiovascular Disease and Global Disease Burden

Cardiovascular diseases rank first as a cause of morbidity in humans worldwide, and sudden death is a common outcome of many of these diseases (coronary atherosclerosis, myocarditis, hypertrophic cardiomyopathy, valve disease, conduction system abnormalities) without discerning age, sex, race or geographical area [1-5]. Sudden death is defined as death that occurs unexpectedly within a 1-hour frame after symptoms start. The causes and mechanisms are unclear, but genetic, structural, functional and environmental factors, among others, are involved, and each will be decisive according to individual characteristics. For example, in children and young individuals, biological and genetic factors are considered determinants; at advanced ages, structural and functional degenerative changes are the main determinants [1-6]. These various possibilities compel the need for studies focusing on these factors; however, clinical research has multiple limitations, including high economic costs, complex logistics, and bioethical considerations, among others. Due to these disadvantages, basic research with animal models is a necessary alternative to investigate this area.

Cardiovascular Research and Animal Models

Multiple animal models have been used for preclinical studies and include invertebrates and vertebrates. The most widely used are Drosophila melanogaster, Caenorhabditis elegans, Xenopus laevis, rodents (rats, mice, rabbits) and birds [7]. Cardiovascular research conducted in these models include embryological and developmental studies, as well as morphological, functional, physiological, pharmacological and toxicological studies [8, 9]. Various published reports have revealed common genetic, molecular, structural and functional cardiovascular patterns for many of the studied species, suggesting the existence of preserved processes and signaling pathways throughout evolution [7]. These signaling pathways common to many species depend on specific expression of genes encoding well-defined molecules that regulate the development of particular cardiovascular structures (Table 1). Genes that are considered key in cardiovascular development and can also be regulated together include GATA, cmlc, nkx, Tbox, and grl, among others [10-12]. In recent decades zebrafish has become an ideal animal model in multiple areas of basic research due to the multiple advantages reported by various authors [7, 13-22]. These advantages include low cost for both acquisition and maintenance, high reproduction, external fertilization and embryogenesis, allowing earlier manipulation of specimens. The life cycle is short, and the period from fertilization to young adult age is completed in three months [13]. Finally, zebrafish has similarity in its genome to humans (approximately 70%, orthologous genes) [14].

 Table 1: Common genes found in zebrafish and cause of sudden death syndrome in humans.

Zebrafish Genes	Protein	Function	Syndromes	References
SCN5A	Nav1.5	I _{Na} , AP, Ph0	BrS	Channelopathies
			LQTS	[1, 2, 6, 38, 41, 72-78]
KCNK1-3-4	TWK-1/2; TASK-1; TRAAK	I _{KP} , AP, Ph0, Ph1, Ph2, Ph3, Ph4	SQTS	
			LQTS	
KCNH2	HERG	I _{Kr} , AP, Ph3	BrS	
			LQTS	
			SQTS	
KCNJ2/12	Kir 2.1/2.2	I _{K1} , AP, Ph3, Ph4	LQTS	
			SQTS	
KCNJ3/5	Kir 3.1/3.4	I _{KAch} , AP, Ph4	LQTS	
KCNQ1	KVLQT1	I _{Ks} , AP, Ph3	LQTS	
			SQTS	
CACNA1C	Cav1.2	ICa,L, AP, Ph2	BrS	

			LQTS SQTS	
RyR2	Ryanodine receptor	Calcium release	ARVD CPVT	Cardiomyopathy [74-76, 79-83]
Sarcomeric genes: cmlc1, amhc, myl7, vmhc, ttn, tnnt2	Troponin T, Titin. myosin light chain,	Contractility	НСМ	

 I_{Na} : Sodium current; I_{Kr} : the fast component of delayed rectifier potassium current; I_{Ks} : the slow component of delayed rectifier potassium current; I_{Kl} : the inward rectifier potassium current; I_{Kr} : the slow component of delayed rectifier potassium current; I_{Kl} : the inward rectifier potassium current; I_{Kr} : leak K+ currents; I_{Ca} , I_{Ca} : I_{Ca} . L-type calcium current; I_{Ca} : AP: Action potential; Ph0: Phase 0; Ph1: Phase 1; Ph2: Phase 2; Ph3: Phase 3; Ph4: Phase 4; BrS: Brugada syndrome; I_{Ca} : Long QT syndrome; SQTS: Short QT syndrome; ARVD: arrhythmogenic right ventricular dysplasia; CPVT: catecholaminergic polymorphic ventricular tachycardia; HCM: hypertrophic cardiomyopathy.

Zebrafish in Cardiovascular Research

As a cardiovascular model, zebrafish has some significant advantages, particularly that the zebrafish heart develops rapidly and is functional at 2 days postfertilization (dpf) and that many of the genes that determine cardiac development have been identified [15, 16]. Additionally, normal and induced cardiovascular morphological and physiological changes can be observed directly using conventional microscopy techniques in zebrafish embryos and larvae due to their transparency [17-22]. Furthermore, during the first week of development, the larvae can survive without circulation because, unlike adult fish, oxygen consumption is independent of cardiac function, as tissues can meet their needs by the simple diffusion of oxygen from the medium [23, 24].

Cardiovascular Development in Zebrafish

The development of the cardiovascular system in zebrafish heart and blood vessels has its origin in the primitive mesoderm where differentiated cells produce a primitive heart disc. The disc progressively evolves into a cardiac cone, from which a tubular structure develops: the cardiac tube. Subsequently, and by specific segmentation and rotation processes, the cardiac tube produces the different well-defined heart chambers: a venous sinus, an atrium, a ventricle and the arterial bulb. These four chambers are arranged in series and are separated by narrowing heart valves that ensure unidirectional blood flow [10, 12, 15, 25-27]. At the same time, hemangioblasts differ from the mesoderm, which produces angioblasts and endothelial cells. At 24 hours postfertilization (hpf), a simple vascular circuit is formed comprising the dorsal aorta and axial vein [28]. From the aorta are derived the intersegmental vessels that will produce different vascular circuits [29].

Heart in Zebrafish: Intrinsic and Extrinsic Activity

Although the cardiac tissue of vertebrates has the properties of excitability, conductivity, contractility and automation, which allow it to work independently, cardiac activity is regulated by the autonomic nervous system (ANS), explaining the dense innervation of the vertebrate heart of vertebrates [30-34]. These characteristics originate in the heart of vertebrates and demonstrate dual cardiac activity-intrinsic activity and extrinsic activity-that is ANS dependent.

I Intrinsic Cardiac Activity

The intrinsic activity is typical of cardiac tissue cells, both noncontractile and contractile cells, which together will determine the

electromechanical activity of the heart. Noncontractile cells determine pacemaker activity, while contractile cells determine cardiac mechanical activity. Both cell types present their resting potentials and action potentials, reflecting the electrical properties of the cell membrane. In zebrafish, the noncontractile cells present in the sinoatrial node generate pacemaker activity by spontaneous depolarization. Two hypotheses have been tested concerning the generation of this activity: the "membrane clock" hypothesis supported by the presence of cyclic nucleotidedependent channels (HCN channels) which favours the ingress of sodium (if); the "calcium clock" hypothesis supported by coupling between intracellular calcium release through active ryanodine receptors and the electrogenic sodium-calcium exchanger (NCX) activation. Some studies have shown that both clocks are coupled in zebrafish heart [35, 36]. Similarly, the action potential of zebrafish cardiac myocytes shares the same characteristics as those in mammals, depends on the interaction of inward and outward currents such as sodium, potassium and calcium and has 5 phases (Table 1). Phase zero is characterized by a rapid ascent that depends on a sodium input current [37]. Phase 1 of repolarization depends on a potassium output current. Phase 2 comprises a plateau due to calcium ingress and potassium output. Phases 3 and 4 depend on potassium currents and cause the return to the resting potential (Table 1). Several studies have shown the presence of counterpart ion channels to those responsible for the intrinsic activity in the heart of mammals [38-411.

II Extrinsic Cardiac Activity

In the neural control of intrinsic cardiac activity, both the sympathetic nervous system (SNS) and parasympathetic nervous system (SNP) participate through two classic neurotransmitters: adrenaline and acetylcholine. Adrenaline works by stimulating cardiac activity through beta-adrenergic metabotropic receptors, while acetylcholine reduces activity by acting through M3 metabotropic receptors. However, some authors have argued that other neurotransmitters, such as GABA and some growth factors, could also be involved in regulating cardiac activity [42-46]. According to current evidence, in zebrafish, an autonomic control is established after the first week of development, implying that, during the first week, cardiac activity depends exclusively on the intrinsic activity. In mammals, asymmetry in cardiac autonomic innervation has been observed, resulting in functional differences. The sympathetic system, through the right stellate ganglion, innervates the sinus node and right atrium, while the left stellate ganglion mainly innervates the left ventricle. The effect of this asymmetry indicates that the activation of the right stellate ganglion increases the heart rate (positive chronotropism) by acting on the sinus node and generates an

anti-arrhythmogenic effect, while the activation of the left stellate ganglion stimulates the myocardium (positive inotropism) and tends to be pro-arrhythmogenic and hypertensive [47].

However, the parasympathetic system regulates cardiac activity through the vagus nerve, which originates in the brain stem in the neurons of the nucleus ambiguus and motor dorsal nucleus of the vagus nerve and its axons extend to intrinsic cardiac nodes located in the wall of the right atrium. Hence, the postganglionic neurons innervate the sinus node [48-50]. Due to this organization, the heart rate varies in time according to the balance between the sympathetic tone and parasympathetic tone. The relationship between alterations in cardiac innervation, particularly cholinergic-adrenergic innervation, and cases of sudden death in arrhythmias, acute myocardial infarction and heart failure has been reported, but the underlying mechanisms involved in these disorders remain unclear [47, 51]. These extrinsic and intrinsic properties present in fish makes zebrafish an ideal model to study intrinsic cardiac activity, neural activity control, its development and maturation and its participation in lethal cardiovascular disorders, such as sudden cardiac death, from the early stages of the life cycle [52-54].

Zebrafish and Heart Disease

The study of sudden cardiac death has been difficult because pathologies of different origins can trigger this phenomenon [4]. The causes of sudden death have ranged from alterations in ion channels and fulminant arrhythmias to metabolic disorders derived from ischaemic phenomena [1, 51]. However, the evidence has not clearly established the physiological mechanisms. In this sense, zebrafish can be an ideal model because models have been characterized to study specific cardiac pathologies. Zebrafish can be used to study the intrinsic properties exclusively in its first week of development, and the determinants of the connection between the heart and autonomic regulation can be evaluated in the second week or the alterations by senescence in the adult zebrafish [35, 39]. In the case of heart disease, models of dilated and hypertrophic heart disease have been established and the relationship between pathology and gene alterations (HAND, NkX2,5, T-box, EYA4 and cat) and/or specific proteins, troponin T, troponin C, titin and sarcomeric proteins [25, 55-61].

The zebrafish has also been a model for studies of the cardiac conduction system, and its alterations are reflected in arrhythmias [62]. Using mutagenic studies of ion channels, such as sodium channels, and optogenetic techniques, arrhythmias have been developed to understand the underlying physiopathological mechanisms [22, 63-66]. Finally, studies investigating cardiac regeneration and myocardial ischaemic disease have used the zebrafish model. Experimental protocols of hypoxia and reperfusion, surgical recession or cryoinjury of the myocardium in the zebrafish ventricle have simulated ischaemia and cardiac reperfusion, a common condition in humans that could also be related to cases of sudden death (Table 1) [67-71].

Conclusion

The wide acceptance of zebrafish in research, the increasingly abundant information on its normal and pathological cardiovascular processes, its short life cycle, its genetic and functional similarities to humans make it an integrated model option to study sudden cardiac death, a multicausal health problem affecting individuals of various ages, races and gender.

Conflicts of Interest

None.

Funding

None.

Consent

Not applicable.

Ethics Committee Clearance

Not Applicable.

REFERENCES

- Fernández Falgueras A, Sarquella Brugada G, Brugada J, Brugada R, Campuzano O (2017) Cardiac Channelopathies and Sudden Death: Recent Clinical and Genetic Advances. *Biology (Basel)* 6: 7. [Crossref]
- Garcia Elias A, Benito B (2018) Ion Channel Disorders and Sudden Cardiac Death. Int J Mol Sci 19: 692. [Crossref]
- Liberthson RR (1996) Sudden Death from Cardiac Causes in Children and Young Adults. N Engl J Med 334: 1039-1044. [Crossref]
- Myerburg RJ (2001) Sudden cardiac death: exploring the limits of our knowledge. J Cardiovasc Electrophysiol 12: 369-381. [Crossref]
- Wong CX, Brown A, Lau DH, Chugh SS, Albert CM et al. (2019)
 Epidemiology of Sudden Cardiac Death: Global and Regional Perspectives. Heart Lung Circ 28: 6-14. [Crossref]
- Chen PS, Choi EK, Zhou S, Lin SF, Chen LS (2010) Cardiac neural remodeling and its role in arrhythmogenesis. *Heart Rhythm* 7: 1512-1513. [Crossref]
- Zaffran S, Frasch M (2002) Early signals in cardiac development. Circ Res 91: 457-469. [Crossref]
- 8. Hasenfuss G (1998) Animal models of human cardiovascular disease, heart failure and hypertrophy. *Cardiovasc Res* 39: 60-76. [Crossref]
- Russell JC, Proctor SD (2006) Small animal models of cardiovascular disease: tools for the study of the roles of metabolic syndrome, dyslipidemia, and atherosclerosis. *Cardiovasc Pathol* 15: 318-330. [Crossref]
- Bakkers J (2011) Zebrafish as a model to study cardiac development and human cardiac disease. Cardiovasc 91: 279-288. [Crossref]
- Reiter JF, Alexander J, Rodaway A, Yelon D, Patient R et al. (1999)
 Gata5 is required for the development of the heart and endoderm in zebrafish. Genes Dev 13: 2983-2995. [Crossref]
- Staudt D, Stainier D (2012) Uncovering the Molecular and Cellular Mechanisms of Heart Development Using the Zebrafish. Annu Rev Genet 46: 397-418. [Crossref]
- Lieschke GJ, Currie PD (2007) Animal models of human disease: zebrafish swim into view. Nat Rev Genet 8: 353-367. [Crossref]
- Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C et al. (2013)
 The zebrafish reference genome sequence and its relationship to the human genome. *Nature* 496: 498-503. [Crossref]
- Hu N, Sedmera D, Yost HJ, Clark EB (2000) Structure and function of the developing zebrafish heart. Anat Rec 260: 148-157. [Crossref]

5

- Stainier DYR, Fishman MC (1994) The zebrafish as a model system to study cardiovascular development. Trends Cardiovasc Med 4: 207-212.
 [Crossref]
- Tanaka T, Oka T, Shimada Y, Umemoto N, Kuroyanagi J et al. (2008)
 Pharmacogenomics of cardiovascular pharmacology:
 pharmacogenomic network of cardiovascular disease models. J
 Pharmacol Sci 107: 8-14. [Crossref]
- Chan PK, Lin CC, Cheng SH (2009) Noninvasive technique for measurement of heartbeat regularity in zebrafish (Danio rerio) embryos. BMC Biotechnol 9: 11. [Crossref]
- Baker K, Warren KS, Yellen G, Fishman MC (1997) Defective "pacemaker" Current (Ih) in a Zebrafish Mutant with a Slow Heart Rate. Proc Natl Acad Sci U S A 94: 4554-4559. [Crossref]
- Chen JN, van Eeden FJ, Warren KS, Chin A, Nüsslein Volhard C et al. (1997) Left-right pattern of cardiac BMP4 may drive asymmetry of the heart in zebrafish. *Development* 124: 4373-4382. [Crossref]
- McGrath P, Li CQ (2008) Zebrafish: a predictive model for assessing drug-induced toxicity. *Drug Discov Today* 13: 394-401. [Crossref]
- Milan DJ, Jones IL, Ellinor PT, MacRae CA (2006) In vivo recording of adult zebrafish electrocardiogram and assessment of drug-induced QT prolongation. Am J PhysiolHeart Circ Physiol 291: H269-H273. [Crossref]
- Warren KS, Fishman MC (1998) "Physiological genomics": mutant screens in zebrafish. Am J Physiol 275: H1-H7. [Crossref]
- Barrionuevo WR, Burggren WW (1999) O2 consumption and heart rate in developing zebrafish (Danio rerio): influence of temperature and ambient O2. Am J Physiol 276: R505-R513. [Crossref]
- Dahme T, Katus HA, Rottbauer W (2009) Fishing for the genetic basis of cardiovascular disease. Dis Model Mech 2: 18-22. [Crossref]
- Holden BJ, Bratt DG, Chico TJA (2011) Molecular control of vascular development in the zebrafish. *Birth Defects Res C Embryo Today* 93: 134-140. [Crossref]
- Liu J, Stainier DYR (2012) Zebrafish in the study of early cardiac development. Circ Res 110: 870-874. [Crossref]
- Serbedzija GN, Flynn E, Willett CE (1999) Zebrafish angiogenesis: A new model for drug screening. Angiogenesis 3: 353-359. [Crossref]
- Childs S, Chen JN, Garrity DM, Fishman MC (2002) Patterning of angiogenesis in the zebrafish embryo. *Development* 129: 973-982.
- Quinn TA, Kohl P, Ravens U (2014) Cardiac mechano-electric coupling research: fifty years of progress and scientific innovation. Prog Biophys Mol Biol 115: 71-75. [Crossref]
- 31. Such L, Rodriguez A, Alberola A, Lopez L, Ruiz R et al. (2002) Intrinsic changes on automatism, conduction, and refractoriness by exercise in isolated rabbit heart. *J Appl Physiol* 92: 225-229. [Crossref]
- Taggart P, Sutton PM (1999) Cardiac mechano-electric feedback in man: clinical relevance. Prog Biophys Mol Biol 71: 139-154. [Crossref]
- Thomas GD (2011) Neural control of the circulation. Adv Physiol Educ
 28-32. [Crossref]
- 34. Valenzuela F, Kabela E (1984) Automatism: an intrinsic property of cardiac tissues. *Arch Inst Cardiol Mex* 54: 601-613. [Crossref]
- Stoyek MR, Croll RP, Smith FM (2015) Intrinsic and extrinsic innervation of the heart in zebrafish (Danio rerio). *J Comp Neurol* 523: 1683-1700. [Crossref]
- Marchant JL, Farrell AP (2019) Membrane and calcium clock mechanisms contribute variably as a function of temperature to setting cardiac pacemaker rate in zebrafish Danio rerio. *J Fish Biol* 95: 1265-1274. [Crossref]

- Grant AO (2009) Cardiac ion channels. Circ Arrhythm Electrophysiol
 185-194. [Crossref]
- Haverinen J, Hassinen M, Dash SN, Vornanen M (2018) Expression of calcium channel transcripts in the zebrafish heart: dominance of T-type channels. J Exp Biol 221: jeb179226. [Crossref]
- MacDonald EA, Stoyek MR, Rose RA, Quinn TA (2017) Intrinsic regulation of sinoatrial node function and the zebrafish as a model of stretch effects on pacemaking. *Prog Biophys Mol Biol* 130: 198-211. [Crossref]
- MacRae CA (2010) Cardiac Arrhythmia: In vivo screening in the zebrafish to overcome complexity in drug discovery. *Expert Opin Drug Discov* 5: 619-632. [Crossref]
- Ravens U (2018) Ionic basis of cardiac electrophysiology in zebrafish compared to human hearts. Prog Biophys Mol Biol 138: 38-44.
 [Crossref]
- 42. Ito T, Hioki H, Nakamura K, Kaneko T, Iino S et al. (2008) Some gamma-motoneurons contain gamma-aminobutyric acid in the rat cervical spinal cord. *Brain Res* 1201: 78-87. [Crossref]
- Llewellyn Smith IJ (2002) GABA in the control of sympathetic preganglionic neurons. Clin Exp Pharmacol Physiol 29: 507-513.
 [Crossref]
- Vargas RA (2016) Effects of GABA, Neural Regulation, and Intrinsic Cardiac Factors on Heart Rate Variability in Zebrafish Larvae.
 Zebrafish 14: 106-117. [Crossref]
- Wang J, Irnaten M, Neff RA, Venkatesan P, Evans C et al. (2001) Synaptic and neurotransmitter activation of cardiac vagal neurons in the nucleus ambiguus. Ann N Y Acad Sci 940: 237-246. [Crossref]
- 46. Wang J, Wang X, Irnaten M, Venkatesan P, Evans C et al. (2003) Endogenous acetylcholine and nicotine activation enhances GABAergic and glycinergic inputs to cardiac vagal neurons. J Neurophysiol 89: 2473-2481. [Crossref]
- Chen PS, Chen LS, Cao JM, Sharifi B, Karagueuzian HS et al. (2001)
 Sympathetic nerve sprouting, electrical remodeling and the mechanisms of sudden cardiac death. *Cardiovasc Res* 50: 409-416.
 [Crossref]
- 48. Armour JA (2008) Potential clinical relevance of the "little brain" on the mammalian heart. Exp Physiol 93: 165-176. [Crossref]
- Mendelowitz D (1996) Firing properties of identified parasympathetic cardiac neurons in nucleus ambiguus. Am J Physiol 271: H2609-H2614. [Crossref]
- Nerbonne JM, Kass RS (2005) Molecular physiology of cardiac repolarization. *Physiol Rev* 85: 1205-1253. [Crossref]
- Pokorný J, Staněk V, Vrána M (2011) Sudden cardiac death thirty years ago and at present. The role of autonomic disturbances in acute myocardial infarction revisited. *Physiol Res* 60: 715-728. [Crossref]
- 52. Billman GE, Huikuri HV, Sacha J, Trimmel K (2015) An introduction to heart rate variability: methodological considerations and clinical applications. *Front Physiol* 6: 55. [Crossref]
- Stauss HM (2003) Heart rate variability. Am J Physiol Regul Integr Comp Physiol 285: R927-R931. [Crossref]
- Stein PK, Bosner MS, Kleiger RE, Conger BM (1994) Heart rate variability: A measure of cardiac autonomic tone. Am Heart J 127: 1376-1381. [Crossref]
- Schönberger J, Wang L, Shin JT, Kim SD, Depreux FFS et al. (2005) Mutation in the transcriptional coactivator EYA4 causes dilated cardiomyopathy and sensorineural hearing loss. *Nat Genet* 37: 418-422. [Crossref]
- Tu S, Chi NC (2012) Zebrafish models in cardiac development and congenital heart birth defects. *Differentiation* 84: 4-16. [Crossref]

Becker JR, Deo RC, Werdich AA, Panàkovà D, Coy S et al. (2011)
 Human cardiomyopathy mutations induce myocyte hyperplasia and activate hypertrophic pathways during cardiogenesis in zebrafish. *Dis Model Mech* 4: 400-410. [Crossref]

- Gerull B, Gramlich M, Atherton J, McNabb M, Trombitás K et al. (2002) Mutations of TTN, encoding the giant muscle filament titin, cause familial dilated cardiomyopathy. *Nat Genet* 30: 201-204. [Crossref]
- Ho YL, Lin YH, Tsai WY, Hsieh FJ, Tsai HJ (2009) Conditional Antisense-Knockdown of Zebrafish Cardiac Troponin C as a New Animal Model for Dilated Cardiomyopathy. Circ J 73: 1691-1697.
- Huang CC, Monte A, Cook JM, Kabir MS, Peterson KP (2013)
 Zebrafish heart failure models for the evaluation of chemical probes and drugs. Assay Drug Dev Technol 11: 561-572. [Crossref]
- Xu X, Meiler SE, Zhong TP, Mohideen M, Crossley DA et al. (2002)
 Cardiomyopathy in zebrafish due to mutation in an alternatively spliced exon of titin. *Nat Genet* 30: 205-209. [Crossref]
- Sedmera D, Reckova M, deAlmeida A, Sedmerova M, Biermann M et al. (2003) Functional and morphological evidence for a ventricular conduction system in zebrafish and Xenopus hearts. Am J Physiol Heart Circ Physiol 284: H1152-H1160. [Crossref]
- Arrenberg AB, Stainier DYR, Baier H, Huisken J (2010) Optogenetic Control of Cardiac Function. Science 330: 971-974. [Crossref]
- Chi NC, Shaw RM, Jungblut B, Huisken J, Ferrer T et al. (2008)
 Genetic and Physiologic Dissection of the Vertebrate Cardiac Conduction System. PLoS Biol 6: e109. [Crossref]
- 65. Huttner IG, Trivedi G, Jacoby A, Mann SA, Vandenberg JI et al. (2013) A transgenic zebrafish model of a human cardiac sodium channel mutation exhibits bradycardia, conduction-system abnormalities and early death. J Mol Cell Cardiol 61: 123-132. [Crossref]
- Chaudhari GH, Chennubhotla KS, Chatti K, Kulkarni P (2013)
 Optimization of the adult zebrafish ECG method for assessment of drug-induced QTc prolongation. *J Pharmacol Toxicol Methods* 67: 115-120. [Crossref]
- 67. Chablais F, Veit J, Rainer G, Jaźwińska A (2011) The zebrafish heart regenerates after cryoinjury-induced myocardial infarction. *BMC Dev Biol* 11: 21. [Crossref]
- González Rosa JM, Martín V, Peralta M, Torres M, Mercader N (2011)
 Extensive scar formation and regression during heart regeneration after cryoinjury in zebrafish. *Development* 138: 1663-1674. [Crossref]
- Parente V, Balasso S, Pompilio G, Verduci L, Colombo GI et al. (2013)
 Hypoxia/reoxygenation cardiac injury and regeneration in zebrafish adult heart. PLoS One 8: e53748. [Crossref]

- Poss KD, Wilson LG, Keating MT (2002) Heart regeneration in zebrafish. Science 298: 2188-2190. [Crossref]
- Schnabel K, Wu CC, Kurth T, Weidinger G (2011) Regeneration of cryoinjury induced necrotic heart lesions in zebrafish is associated with epicardial activation and cardiomyocyte proliferation. *PLoS One* 6: e18503. [Crossref]
- Arnaout R, Ferrer T, Huisken J, Spitzer K, Stainier DYR et al. (2007)
 Zebrafish model for human long QT syndrome. *Proc Natl Acad Sci U S A* 104: 11316-11321. [Crossref]
- Brown DR, Samsa LA, Qian L, Liu J (2016) Advances in the Study of Heart Development and Disease Using Zebrafish. J Cardiovasc Dev Dis 3: 13. [Crossref]
- Kauferstein S, Kiehne N, Neumann T, Pitschner H-F, Bratzke H (2009)
 Cardiac Gene Defects Can Cause Sudden Cardiac Death in Young
 People. Dtsch Arztebl Int 106: 41-47. [Crossref]
- Kimmel CB (1989) Genetics and early development of zebrafish. *Trends Genet* 5: 283-288. [Crossref]
- Perrin MJ, Gollob MH (2012) The Genetics of Cardiac Disease Associated with Sudden Cardiac Death: A Paper from the 2011 William Beaumont Hospital Symposium on Molecular Pathology. J Mol Diagn 14: 424-436. [Crossref]
- Skarsfeldt MA, Bomholtz SH, Lundegaard PR, Lopez-Izquierdo A, Tristani-Firouzi M et al. (2018) Atrium-specific ion channels in the zebrafish - A role of IKACh in atrial repolarization. *Acta Physiol (Oxf)* 223: e13049. [Crossref]
- Verkerk AO, Remme CA (2012) Zebrafish: a novel research tool for cardiac (patho) electrophysiology and ion channel disorders. Front Physiol 3: 255. [Crossref]
- Collins MM, Ahlberg G, Hansen CV, Guenther S, Marín-Juez R et al. (2019) Early sarcomere and metabolic defects in a zebrafish pitx2c cardiac arrhythmia model. *Proc Natl Acad Sci U S A* 116: 24115-24121. [Crossref]
- Lin E, Shafaattalab S, Gill J, Al-Zeer B, Craig C et al. (2020) Physiological phenotyping of the adult zebrafish heart. *Mar Genomics* 49: 100701. [Crossref]
- 81. Shi X, Chen R, Zhang Y, Yun J, Brand-Arzamendi K et al. (2018) Zebrafish heart failure models: opportunities and challenges. *Amino Acids* 50: 787-798. [Crossref]
- 82. Yan J, Li H, Bu H, Jiao K, Zhang AX et al. (2020) Aging-associated sinus arrest and sick sinus syndrome in adult zebrafish. *PLoS One* 15: e0232457. [Crossref]
- Zhu X-Y, Wu S-Q, Guo S-Y, Yang H, Xia B et al. (2018) A Zebrafish Heart Failure Model for Assessing Therapeutic Agents. Zebrafish 15: 243-253. [Crossref]